

Practical Strategies for Preserving Good Cognitive Function into Old Age

Mark F. McCarty, Catalytic Longevity, markfmccarty@gmail.com

As people age, their cognitive function is threatened by the normal aging process, as well as increasing risk for stroke and a range of inflammatory neurodegenerative conditions, most notably Alzheimer's disease (AD), but also including frontotemporal dementia, Parkinson's disease and others. Our understanding of the origins of these disorders, and of the most definitive ways to prevent, postpone, or treat them is still at a rudimentary stage – which is understandable, in light of the phenomenal complexity of cellular biology and the human brain. Nonetheless, I submit that, based on current evidence, a range of practical prudent strategies can be defined that seem likely to aid preservation of cognitive function during aging and that, in any case, appear consistent with promotion of optimal health:

Prudent Diet and Lifestyle – Epidemiological studies strongly suggest that a diet pattern characterized as “Mediterranean” or plant-based – low in saturated fats, and rich in fruits and vegetables – can slow cognitive decline during aging while reducing risk for AD and other dementing disorders.¹⁻⁵ With respect to stroke risk, moderation in dietary salt and a high intake of potassium-rich whole foods appears to be notably protective. Regular exercise training also emerges as globally protective – slowing the rate of age-related cognitive decline, and reducing risk for AD and stroke. And – consistent with the adage “use it or lose it” – some studies point to regular mental exercise as beneficial for cognitive preservation.

The risk for cognitive decline associated with diets high in saturated fats has recently been traced to the fact that these fats, unlike unsaturated fats, can give rise to the pro-inflammatory compound ceramide, which in turn promotes the pro-inflammatory activation of brain glial cells – astrocytes and microglia.⁶⁻¹¹ Saturated fats can also activate microglia by stimulating their “toll-like” receptors.^{11, 12} These cells function in various ways to support healthy brain function. In particular, they protect the brain from invasion by infective agents, for which reason they are capable of generating a wide range pro-inflammatory hormones as well as oxidant chemicals such as peroxynitrite. A notable downside of this is that, in excess, these hormones and oxidants can perturb the proper function of brain neurons; they can impede the efficiency of a process known as “long-term potentiation” required for new memory formation, can amplify the production and potentiate the effects of the amyloid beta protein that is a key driver of neuronal dysfunction and death in AD, and can even kill neurons.^{9, 13-17} As people age, there is a natural tendency for brain glial cells to become more pro-inflammatory – either they are chronically active in producing pro-inflammatory hormones (notably interleukin-1), or are on hair-trigger alert, highly sensitive to activating stimuli. This phenomenon is thought to be largely responsible for the modest loss of cognitive function normally associated with “healthy aging”, and also has the potential to exacerbate incipient neurodegenerative disorders such as AD or Parkinson's.^{15, 18, 19} Hence, aging is associated with chronic low-grade inflammation of the brain – some might call it a “brain on fire” syndrome. Eating a diet rich in saturated fat is like throwing fuel into this fire. As we will see, many of the phytonutrients or drugs discussed below with potential for slowing age-related cognitive decline can suppress the tendency of brain glial cells to become inflamed - and hence have an opposite effect to that of saturated fat.

Dietary intakes of sodium and potassium appear to be a key determinant of cerebrovascular health and stroke risk. Hypertension is of course a major risk factor for both types of stroke – ischemic (triggered by

blood clots) and hemorrhagic. In societies that don't salt their food, the most common type of hypertension – the so-called “essential” form – is virtually non-existent; moreover, there is limited evidence that both stroke and dementia are quite rare in these societies.^{20, 21} Few Western studies have examined the association of salt intake with risk for dementia, but one recent study found that relatively low dietary sodium intake in elderly people was associated with a lower rate of cognitive decline over the subsequent three years.²² Of related interest are reports that memory is superior and risk for dementia is notably lower in hypertensives who use potassium-sparing diuretics; these drugs cause urinary sodium loss while preventing loss of potassium.^{23, 24} In societies that do salt their food, many but not all hypertensives are said to be “salt-sensitive”, meaning that a reduction in dietary salt can bring their blood pressure down a bit. A high intake of potassium tends to oppose the pro-hypertensive impact of dietary salt, and clinical studies show that an important increase in dietary potassium tends to lower blood pressure.²⁵ But what is at least as important is that, at any given level of blood pressure, a higher intake of potassium is associated with a lower risk for stroke²⁶⁻³³ – likely because a modest increase in blood potassium levels has a direct antioxidant/anti-inflammatory impact on the endothelial lining of the cerebrovascular arteries.³⁴⁻³⁶ A lower dietary sodium/potassium ratio likewise has a favorable impact on risk for heart attack.³⁷⁻³⁹

In a very impressive recent controlled clinical study, veterans in a Chinese retirement home were randomized to receive either their ordinary diet, or a very similar diet in which a salt substitute (half sodium chloride, half potassium chloride) replaced ordinary salt in cooking and at the table.⁴⁰ Use of the salt substitute led to only a modest reduction in salt intake (owing to the use of some pre-salted ingredients), but potassium intake went up by about 76% in this group. During a follow-up of about two and a half years, the death rate from heart attack, stroke, or heart failure was 40% lower in those getting the salt substitute! Moreover, a population-wide reduction of the dietary sodium-potassium ratio in Finland appears to have made an important contribution to the dramatic reduction in heart attack and stroke rates observed in that country over the last two decades.⁴¹ You can notably increase the potassium content of your diet by emphasizing whole foods (added sugars, fats, and distilled liquors are virtually devoid of potassium, and refined grains are a poor source); some fruits, such as bananas, are exceptionally rich in potassium, as are most tubers. When you use salt in cooking or at the table, using a salt substitute in which potassium chloride partially replaces sodium chloride is a shrewd move.

In many epidemiological studies, regular exercise emerges as protective with respect to risk for stroke, age-related cognitive decline, and dementia.⁴²⁻⁴⁹ Part of this protection no doubt stems from the favorable impact of exercise on cerebrovascular function and hypertension.⁵⁰ But for reasons that remain unclear, exercise also provokes the brain to produce increased levels of growth factors – so-called neurotrophic factors such as brain-derived neurotrophic factor (BDNF) – that aid the survival and effective function of neurons.⁵¹⁻⁵⁴ There is a tendency for the hippocampus – a portion of the brain cortex vital for new memory formation – to shrink a bit during aging; a recent controlled study found that one year of aerobic exercise in adults over 55 was associated with a modest but significant increase in hippocampal volume.⁵⁵ In rodent models of Alzheimer's disease – mice bioengineered to overproduce the amyloid beta protein that drives cognitive dysfunction and neuronal death in this syndrome – voluntary exercise favorably affects their cognitive function and in some instances lowers the deposition of amyloid beta aggregates (plaque) in the brain.⁵⁶⁻⁶¹ In normal aging mice, exercise training appears to alter the function of brain microglia so that they are less pro-inflammatory and more supportive of brain health.⁶² In short,

all signs point to a lifelong habit of exercise as being notably protective for cognitive health during aging, and even suggest that taking up exercise late in life can be beneficial in this regard.^{44, 63}

One of the “side effects” of an exercise habit is that it helps you to control your weight; a Mediterranean or plant-based diet emphasizing whole foods can also be quite helpful in this regard.⁶⁴ People who are overweight in midlife – particularly if they have a preferential accumulation of body fat in their belly, known as android obesity – are at notably higher risk for AD and age-related cognitive decline in later years.⁶⁵ Analogously, the metabolic abnormalities typically associated with android obesity, known as the “metabolic syndrome” (insulin resistance, hypertension, high triglycerides, low HDL) are also mid-life risk factors for late-life dementia.⁶⁶⁻⁷⁰ These findings serve to emphasize the fact that AD is a chronic disorder that develops gradually over decades, only becoming symptomatic after a number of years. Surprisingly, overweight does not consistently emerge as a risk factor for AD among elderly people, and associations between metabolic syndrome and AD in the elderly are sometimes inconsistent.^{65, 71} Conceivably, this may simply reflect the fact that the weight loss which is a typical consequence of AD^{63, 72} may often precede diagnosis of the syndrome; if that is the case, then correction of metabolic syndrome or obesity in the elderly might still have utility for diminishing AD risk. In this regard, diabetes, typically associated with overweight, is indeed a major risk factor for cognitive decline and AD in the elderly.^{66, 73, 74} Adopting healthful habits that tend to promote a leaner physique – eating a prudent diet and exercising regularly and effectively – is probably a smart idea at any age.

In the context of diet, it may be appropriate to mention coffee and caffeine. The “mental-energizing” effects of caffeine reflect its ability to inhibit certain cellular receptors in the brain which ordinarily are activated by the natural metabolite adenosine. Via interaction with the A2A form of adenosine receptor, caffeine has the potential to exert anti-inflammatory effects on the brain, dampening the activation of microglia, supporting the protective function of astrocytes, decreasing production of beta amyloid, and warding off some of the adverse effects of beta amyloid on neuronal synapses and cognitive function.⁷⁵⁻⁸² Indeed, most though not all recent epidemiological studies find that people who have made heavy use of coffee for a number of years are at decidedly lower risk for cognitive decline and AD as they age.⁸³⁻⁸⁹ In one provocative Finnish study, people who drank 3-5 cups of coffee daily in midlife were about *two-thirds* less likely to develop dementia as they aged.⁸⁶ This one study probably overstates the true benefit of caffeine – some studies find no protection^{90, 91} - but the balance of evidence clearly points to a notable protective effect. Whether adopting a caffeine habit at an advanced age can be protective in this regard has not yet been studied. However, one recent prospective study examined blood caffeine levels in elderly subjects with mild cognitive impairment, and found that caffeine level dose-dependently predicted risk for development of dementia over the next 4 years – none of those in the highest bracket of blood caffeine became demented over this time.⁹² Other recent epidemiology on coffee points to heavy coffee drinkers as being at reduced risk for mortality from a host of disorders, including heart disease, stroke, diabetes, infections, injuries, and accidents; some of this protection might stem, not just from caffeine, but from the phytochemical antioxidants in coffee.⁹³ In light of current evidence, it may be smart to drink about as much coffee as you can, consistent with getting appropriate restful sleep and the concurrence of your physician. If drinking coffee in the evening or even afternoon impairs your sleep, you can limit your use to the morning. (Some decades ago, caffeine and coffee developed a reputation as a potential health *risk*, owing to the fact that, in people who are coffee “virgins”, a dose of caffeine provokes an increase in heart rate and blood pressure. But if people persist in drinking coffee regularly, this reaction tends to dampen and disappear over several weeks; that’s why coffee doesn’t harm regular coffee drinkers.)

Caffeine's particular merit for cognitive health may be its ability to block the adverse impact of amyloid beta – the toxic protein that is a key driver of neuronal death and dysfunction in AD – on cellular uptake of the neurotransmitter glutamate. Glutamate is major neurotransmitter in the brain, including those parts of the brain responsible for memory formation. After its extracellular release at synapses is triggered by a neural impulse, it must be rapidly removed from the extracellular space by cellular uptake; glial cells known as astrocytes are primarily responsible for this uptake. But amyloid beta suppresses the ability of astrocytes to take up glutamate, and there is recent evidence that it does this via stimulation of astrocyte A2A adenosine receptor activity.^{94, 95} Failure to rapidly remove glutamate after a neural impulse results in chronic low-grade stimulation of a specific type of neuronal glutamate receptor (so-called extrasynaptic NMDA receptors), and this is now suspected to be a major mechanism whereby amyloid beta impairs neuronal function and ultimately can cause neuronal death.⁹⁶⁻⁹⁸ By blocking A2A receptors, caffeine negates the adverse impact of amyloid beta on astrocyte glutamate uptake, and hence may strike at the root of a key mechanism whereby amyloid beta damages neurons.

One of the long-chain omega-3 fatty acids found typically in fish oil – docosahexaenoic acid (DHA) – is a prominent structural component of neuronal cell membranes, both in the brain and retina. The process of establishing new memories requires rapid synthesis of new connections between neurons – a process known as synaptogenesis – and this in turn requires efficient production of new neuronal membranes. As DHA is a major component of these membranes, a ready availability of DHA aids the efficiency of this process.⁹⁹ Hence, it is not surprising that DHA-enriched diets have shown favorable effects on cognitive function in rodent studies – though not always in AD model mice.¹⁰⁰⁻¹⁰⁷ Epidemiologically, elderly people with relatively high blood levels of DHA or high fish intakes tend to enjoy superior cognitive function and lower risk for dementia, and good DHA status may predict a lower rate of cognitive decline.^{1, 100, 108-113} In one recent controlled clinical study, volunteers over 55 performed better on certain tests of learning and memory after they had received 900 mg of supplemental DHA daily.¹¹⁴ However, supplemental DHA has not been found to slow cognitive decline in AD.^{109, 115} You can insure good DHA status if you eat oily fish regularly, or if you take a fish oil supplement rich in omega-3. Krill oil has also been introduced recently as a supplemental source of DHA, and vegans can obtain DHA supplements derived from marine algae (albeit these are currently far more expensive than fish oil per unit DHA). Although humans inefficiently convert the main omega-3 fat found in plant products – alpha-linolenic acid – to DHA, a novel type of soybean oil rich in stearidonic acid, an omega-3 more readily converted to DHA, should soon be on the market.¹¹⁶

Another nutrient that is emerging as especially beneficial for cognitive health is vitamin D. When microglia become inflamed, their capacity to convert circulating vitamin D to its active hormonal form goes up, and the resulting increase in microglial vitamin D activity acts as a restraint on microglial inflammation.¹¹⁷⁻¹¹⁹ As noted, interleukin-1 (IL-1) is a key hormone produced by inflamed microglia that can impair the efficiency of the long-term potentiation required for memory formation in the hippocampus; in aging rats treated with vitamin D, hippocampal levels of IL-1 declined significantly.¹²⁰ Recent epidemiological studies are finding that, among aging people, poor vitamin D status tends to correlate with poorer cognitive status, and also is associated with increased risk for cognitive decline and development of dementia.¹²¹⁻¹²⁸ Poor vitamin D status is also being linked to increased risk for vascular disease, diabetes, and certain prominent types of cancer and autoimmunity.¹²⁹ Although limited amounts of vitamin D can be obtained from food, most people synthesize most of their vitamin D via exposure of their skin to *uv* light; however, the *uv* content of sunlight tends to be quite low at northern latitudes during

the winter. Hence, a growing number of health authorities are recommending a daily supplemental intake of vitamin D, in the range of 2,000-8,000 IU daily, as the most practical way to insure good vitamin D status year- round; this appears to be quite safe, as our own bodies can manufacture over 10,000 IU daily.¹³⁰

Cerebral Antioxidants – There is a growing consensus that cerebral oxidative stress plays a key role in the induction of age-related cognitive decline, AD, Parkinson’s disease, Lewy body dementia, and stroke. Cerebral oxidative stress arises primarily from overactivation of an enzyme complex known as NADPH oxidase, and also from mitochondria (the “power plants” inside our cells that manufacture the energy catalyst ATP) that have been structurally disrupted. Both NADPH oxidase and damaged mitochondria make superoxide, which in turn is converted to downstream compounds – hydrogen peroxide, hydroxyl radical, and peroxynitrite – which are unstable and, if not promptly detoxified by antioxidant enzymes or small molecule antioxidants such as glutathione, can cause severe structural damage to cells and/or activate pro-inflammatory cellular signaling pathways. In microglia, activation of NADPH oxidase is a key driver of pro-inflammatory behavior, and can damage neighboring cells by giving rise to the oxidant compound peroxynitrite.^{131, 132} Oxidative damage of neurons by activated microglia is thought to be a prime mediator of Parkinson’s disease, and may also contribute to neuronal damage in AD and Lewy body dementia.¹³³⁻¹³⁸ Neurons also can express NADPH oxidase, which can be stimulated by amyloid beta^{139, 140} – but activation of mitochondrial superoxide production appears to be a primary way in which toxic amyloid beta promotes neuronal dysfunction and death in AD.¹⁴¹⁻¹⁴⁴ There is recent evidence that, in mice, chronic activation of NADPH oxidase in a special subset of brain neurons (GABAergic interneurons) promotes the age-related die-off of these neurons, which in turn contributes to age-related cognitive dysfunction.^{145, 146} The amyloid beta-driven inflammatory process in AD is often associated with local reductions in blood flow attributable to amyloid beta-mediated activation of NADPH oxidase in the cerebral microvasculature; this process, known as cerebral amyloid angiopathy, is thought to exacerbate cognitive decline in AD.^{147, 148} Activation of NADPH oxidase in the cerebral vasculature may also play a key pathogenic role in the induction of stroke and in the resulting damage to the brain.¹⁴⁹⁻¹⁵³ Hence, effective control of cerebral oxidative stress may have potential for stemming age-related cognitive decline, reducing stroke risk, and slowing the onset or progression of dementing neurodegenerative disorders.

Two phytonutrients stand out as having particularly outstanding potential for controlling cerebral oxidative stress – phycocyanobilin (PhyCB), richly supplied by the food microalga spirulina, and astaxanthin, currently produced for supplemental use from another type of microalga. Bilirubin, a compound produced naturally in our bodies, has remarkable antioxidant activity because it can directly inhibit several forms of the NADPH oxidase complex.¹⁵⁴⁻¹⁵⁶ (Indeed, when cells are under oxidative stress, this provokes the synthesis of an enzyme, heme oxygenase-1, which generates bilirubin within the cell, thereby providing feedback control of the oxidative stress.¹⁵⁷) Although there are no natural concentrated sources of bilirubin for supplemental use, it is a great stroke of luck that, within our cells, PhyCB can be rapidly converted to a compound almost identical in structure to bilirubin which likewise can potently inhibit NADPH oxidase.¹⁵⁸⁻¹⁶⁰ This provides a very credible explanation for the fact that, in rodent studies, oral administration of whole spirulina or of protein-bound PhyCB (known as phycocyanin) has shown protective effects in a great many models of inflammation.^{159, 161} Indeed, PhyCB may have the potential to provide protection from a vast number of health disorders in which oxidative stress generated via NADPH oxidase plays a mediating or exacerbating role, including vascular diseases, diabetic

complications, and autoimmune conditions.¹⁵⁹ Moreover, there is good reason to suspect that orally administered PhyCB has access to the brain, as oral administration of spirulina or of phycocyanin has been found to be protective in rodent models of Parkinson's disease and epilepsy.¹⁶²⁻¹⁶⁴ More pertinently, in SAMP8 mice, a "senescence-accelerated" mouse strain prone to early memory decline and increased brain production of amyloid beta, three months of spirulina feeding prevents loss of memory function in 6-month-old mice, while notably quelling brain oxidative stress.¹⁶³ Another recent study shows that spirulina feeding preserves the function of neuronal stem cells (cells capable of generating new neurons) in rats whose brains had been injected with the powerful pro-inflammatory compound lipopolysaccharide.¹⁶³ And exposure of microglial cells to phycocyanin opposes their inflammatory activation – as had been predicted.^{165, 166} Also of interest is a study showing that, if gerbils are fed phycocyanin for several days before they are subjected to a simulated stroke, the degree of brain damage that results is lessened.¹⁶⁷

Hence, although clinical assessments of spirulina's cerebroprotective potential are currently lacking, rodent studies and cell culture studies suggest it may have profound potential in this regard. Presuming that humans absorb and metabolize PhyCB in a manner comparable to rodents, it has been estimated that a daily intake of 15-30 g spirulina daily (1-2 tablespoons) could be expected to replicate the protective antioxidant effects observed in rodents.¹⁵⁹ Consuming this much spirulina can be a daunting challenge, as spirulina tastes rather bad and smells even worse, but functional foods in which the flavor of spirulina is masked are being developed, and it should be possible someday for biotechnologists to extract the PhyCB from spirulina and use it in nutraceuticals.

As we have noted, damaged mitochondria are thought to be a major source of pathogenic oxidative stress in AD. Astaxanthin, a fat-soluble membrane antioxidant similar in structure but more potent in antioxidant activity than the carotenoid phytonutrients lutein and zeaxanthin – and many times more active than vitamin E – appears to have considerable potential for protecting cellular membranes, and most particularly the membranes of mitochondria, which are under continual threat from oxidative stress.¹⁶⁸⁻¹⁷² Although healthy mitochondria produce superoxide at a low, tolerable rate, oxidative damage to mitochondrial membranes can turn mitochondria into avid sources of superoxide – which in turn provokes more damage to the mitochondrial membranes. Astaxanthin has the potential to block this vicious cycle by preserving membrane structure. This is likely to be the mechanism whereby astaxanthin is protective in so-called "ischemia-reperfusion" damage, which plays a major role in tissue damage in both heart attack and stroke.^{173, 174} Unfortunately, astaxanthin has not yet been tested in rodent models of AD or Parkinson's – but it will be surprising if it doesn't show some protection in these models; one recent review cites its neuroprotective potential and calls it "a potent candidate for brain food".¹⁷⁵ There are however reports that astaxanthin can exert an anti-inflammatory effect on microglial cells in culture.^{176, 177} This is perhaps a bit surprising, since NADPH oxidase is the chief source of oxidative stress in these cells – but perhaps this reflects a role for metabolites of oxidatively-damaged membranes in the induction of inflammation in microglia. Although astaxanthin, like spirulina, has so far received little clinical evaluation, initial studies suggest that supplemental intakes of 4-20 mg daily can have a significant antioxidant impact and clinical utility.¹⁷⁸⁻¹⁸² In light of clinical evidence that vitamin E may modestly slow cognitive decline in AD, the utility of astaxanthin in this regard certainly merits evaluation.¹⁸³

An alternative approach to controlling cerebral oxidative stress is to administer natural compounds which can provoke increased synthesis of antioxidant enzymes in the brain cells. Two categories of compounds have this potential – so-called “phase 2 inducers”, and the hormone melatonin. The phase 2 inducers include a vast number of phytochemicals, many of which are found in common foods – sulforaphane in cruciferous vegetables, EGCG in green tea, sulfur compounds found in garlic, etc. – and notably the nutritional cofactor lipoic acid, which has shown neuroprotective activity in a number of rodent studies and in human diabetic neuropathy.¹⁸⁴⁻¹⁹¹ (The cerebroprotective benefits of some of the polyphenols discussed in the next section may stem in part from phase 2 induction.) Phase 2 inducers, by boosting the level and promoting the nuclear uptake of the intracellular protein nrf2, cause cells to increase their synthesis of a large number of antioxidant enzymes, and to also increase their production of the crucial intracellular antioxidant glutathione.¹⁹²⁻¹⁹⁵ None too surprisingly, there is evidence that boosting nrf2 expression in the brains of AD model mice (dementia-prone mice genetically engineered to produce increased levels of amyloid beta) improves their cognitive function.¹⁹⁶ Among the phase 2 inducers, lipoic acid is of particular interest, as it quite clearly has access to the brain; dietary supplementation with lipoic acid is reported to improve cognitive function in AD model mice.¹⁹⁷⁻¹⁹⁹ The daily dose of lipoic acid shown to be effective in treating diabetic nerve damage are in the 600-1800 mg daily range; likely such doses would also provide some protection to the brain.¹⁹⁷ A couple of small pilot clinical studies with lipoic acid suggest that it may indeed slow the onset of dementia in early AD.^{200, 201}

Melatonin is a hormone produced by the pineal gland at the base of the brain, and also synthesized by some cells. It is released during the night, and helps to coordinate the body’s biorhythms; this secretion tends to diminish with increasing age, which is why supplemental melatonin may be most helpful in the elderly. Melatonin has an impact similar to phase 2 inducers on cellular antioxidant enzymes and glutathione; however, its effects are not dependent on activation of nrf2.²⁰² Like lipoic acid, melatonin aids cognitive function in AD mouse models; in cell cultures, it protects neurons from amyloid beta-mediated damage, and lessens the proinflammatory impact of amyloid beta on microglia.²⁰³⁻²¹⁰ Fortunately, melatonin is absorbed efficiently when administered orally. When used clinically, melatonin is administered at bedtime (to mimic the normal rhythm of its physiological production), usually in a dose of 3-20 mg. Some people may find they are a bit groggy the next morning if they take too high a dose. Initial clinical evaluation of melatonin in patients with AD suggests that it may be beneficial for a syndrome known as “sundowning”, in which patients become agitated with the onset of evening; this likely reflects its ability to restore normal biorhythms.²¹¹

As noted, one key way in the phase 2 inducers and melatonin increase the antioxidant defenses of neurons is to boost production of the versatile intracellular antioxidant glutathione; they do this by increasing the expression of the enzyme whose activity is rate-limiting for its production. Glutathione is synthesized within cells from the amino acids cysteine, glutamic acid, and glycine; cysteine is usually the least available of these amino acids, and hence its concentration determines the speed with which glutathione can be synthesized. Hence, supplementation which increases intracellular cysteine levels can increase cellular levels of glutathione. Although high oral doses of cysteine per se cause GI upset, supplementation with the compounds N-acetylcysteine (NAC) or cystine is far better tolerated, and accomplishes the purpose of boosting intracellular cysteine - and hence glutathione.²¹²⁻²¹⁵ Protective benefits of NAC supplementation have been reported in rodent models of Parkinson’s disease and AD.²¹⁶⁻

223

Lewy body dementia (DLB), although it so far has received little publicity, is thought to be responsible for 20% or more of clinical dementia. Its origins appear to be similar to those of Parkinson's disease (PD), because in each of these disorders intraneuronal aggregates of the protein alpha-synuclein, known as Lewy bodies, are believed to contribute to neuronal dysfunction and death.¹³⁸ DLB begins in cortical regions of the brain, whereas PD arises in the substantia nigra, rich in neurons that make the neurotransmitter dopamine. Curiously, patients with DLB often eventually develop Parkinsonian symptoms, whereas PD patients are prone to develop dementia associated with cortical Lewy bodies; it is reasonable to suspect that these disorders are, from a mechanistic standpoint, very similar, but differing in where they first arise in the brain. Microglial activation and microglial production of the vicious oxidant peroxynitrite appear to be key mediators of the destruction of dopaminergic neurons that characterizes PD; not unlikely, these factors may also drive DLB.^{138, 224-228} Peroxynitrite is toxic, in part, because it promotes the aggregation of alpha-synuclein and the formation of Lewy bodies.²²⁹⁻²³² Microglial production of peroxynitrite, which arises from a spontaneous reaction from the compounds superoxide and nitric oxide, can be suppressed by inhibiting microglial superoxide production with PhyCB – likely accounting for the favorable impact of spirulina in rodent models of PD.^{162, 164} Furthermore, certain natural antioxidants can provide protection from the pathogenic effects of peroxynitrite – notably the intracellular antioxidant glutathione.²³³ Hence, phase 2 inducers, melatonin, and cysteine sources (NAC, cystine) may be useful for controlling PD and DLB – consistent with results in rodent studies.^{216-221, 234-237}

But the natural metabolite uric acid (urate) also has important potential for detoxifying peroxynitrite;²³⁸ this may explain why people whose urate levels are naturally high are at significantly decreased risk for PD, or have a slower progression of their disease if they already have this disorder.²³⁹ Unfortunately, the impact of urate on risk for DLB has not yet been studied; however, patients with PD may be at lower risk for subsequent dementia if their urate levels are high^{240, 241} – a finding consistent with the possibility that urate provides protection from DLB as well. Supplementation with the purine compound inosine can raise urate levels, and urate supplementation is now being studied in the treatment of multiple sclerosis, another neurodegenerative disorder in which peroxynitrite is suspected to play a key pathogenic role.²⁴²⁻²⁴⁴ However, supplementation with inosine is not entirely innocuous, because excessive blood and tissue levels of urate can lead to the deposition of uric acid crystals in the body's tissues and kidneys, resulting in the agonizing arthritic disorder gout. Gouty arthritis can be prevented during inosine supplementation if the dose is titrated such that serum urate levels are kept within the high-normal range 6-9 mg/dL; in the multiple sclerosis clinical trials, physicians start with an intake of 1 g inosine daily, and gradually increase the dose to as much as 3 g daily, in an effort to keep urate in a high but safe target range.²⁴⁴ However, even if careful dose adjustment succeeds in preventing gouty arthritis, patients supplemented with inosine are at increased risk for uric acid kidney stones.²⁴⁴ This risk can be minimized if the urine is kept alkaline (as urate tends to crystallize under acidic conditions); a moderate-protein plant-based diet high in potassium helps to maintain an alkaline urine, and supplementation with potassium bicarbonate or potassium citrate can also be employed for this purpose. Because of its logistical difficulty, inosine supplementation can't be recommended for primary prevention of neurodegenerative disorders, but with the supervision of a cooperative physician, it may be worthy of consideration in patients who are developing PD or DLB. Measures which dampen microglial activation (as discussed below), boost neuronal glutathione levels, and protect mitochondrial membranes (such as astaxanthin²⁴⁵⁻²⁴⁷) might also prove useful in these disorders.

Acetyl-L-carnitine (ALC) is often thought of as a brain antioxidant, as it does exert antioxidant effects on the brain in certain circumstances, but the basis of its antioxidant activity remains rather obscure.²⁴⁸⁻²⁵⁰ Speculation centers on the possibility that it may reduce oxidant production by damaged mitochondria. Dr. Bruce Ames and colleagues have shown that, in aging rodents, supplementing with the combination of lipoic acid and ALC aids the production of new mitochondria in various tissues, including the brain.²⁵⁰⁻²⁵³ Since new mitochondria are less prone to generate oxidative stress than damaged old mitochondria, perhaps this helps to explain how ALC acts as an antioxidant. The ability of lipoic acid/ALC to boost brain production of mitochondria is of particular interest in light of the fact that amyloid beta provokes mitochondrial damage in AD. ALC also has the potential to promote production of the neurotransmitter acetylcholine, as discussed below.²⁵⁴⁻²⁵⁶ Regardless of how ALC influences brain function, a great many clinical trials have conclusively demonstrated that, in daily doses of 2-3 grams, ALC has a beneficial impact on cognitive function in elderly people with minimal cognitive impairment or early AD.²⁵⁷ Despite this, there appear to be few published studies evaluating its impact in mouse AD models.

Taurine is a vitamin-like cofactor that has a special antioxidant function (quenching the hypochlorous acid produced by activated macrophages), but it also regulates membrane function and calcium metabolism in a range of tissues. Neurons are among the cells that accumulate taurine. As you will recall, there is recent evidence that age-related loss of GABAergic interneurons plays a role in normal cognitive aging. A New York research group has recently reported that, when mice were supplemented with taurine for 8 months beginning at 8 months of age, the cognitive decline typically seen in 16-month-old mice was prevented; this effect was thought to reflect protection of a subset of GABAergic interneurons.^{258, 259} Aside from this provocative report, researchers have so far shown little interest in exploring the long-term effects of taurine on cognitive function. In any case, taurine is inexpensive and quite safe, and there is considerable reason to believe that optimal taurine status may be protective for vascular health. Supplemental intakes in the range of 1-6 g daily are feasible, as taurine is highly soluble in fluid and has no flavor (in fact, it is a key ingredient of Red Bull™!)

Controlling hypertension is of course well known to be beneficial for stroke prevention, and this often entails use of anti-hypertensive drugs. There is recent evidence that brain-permeable drugs which antagonize the function of the pro-hypertensive factor angiotensin II (drugs known as ACE inhibitors and type 1 angiotensin receptor antagonists) can exert an anti-inflammatory/antioxidant effect on the brain, and moreover have a beneficial impact on mouse AD models.²⁶⁰⁻²⁶⁵ This likely is pertinent to humans as well.²⁶⁶⁻²⁶⁸ In one recent epidemiological study, in which the age-related cognitive decline of treated hypertensives was assessed, those treated with ACE inhibitors capable of entering the brain showed a 65% lesser decline over 6 years than the group as a whole – whereas those receiving ACE inhibitors that did not have access to the brain declined at a slightly higher rate than the whole group.²⁶⁹ Other recent epidemiology points to lower risk for dementia in users of angiotensin receptor antagonists.²⁷⁰ In salt-sensitive rats, a high-salt diet has recently been shown to boost hippocampal levels of angiotensin II and impair cognitive function; concurrent treatment with an angiotensin receptor antagonist alleviated the cognitive impairment.²⁷¹ This may be germane to previously cited evidence that a relatively low dietary salt intake is associated with lesser risk for cognitive decline. ACE inhibitors with the ability to cross the blood-brain barrier include captopril, fosinopril, lisinopril, perindopril, ramipril, andtrandolapril.²⁶⁹ Angiotensin receptor antagonist drugs have access to the brain, and seem likely to be comparably protective for brain function.

Anti-inflammatory Polyphenols – A vast and growing research literature focuses on the potential of dietary polyphenols for aiding preservation of cognitive function during aging . Polyphenols, found in a wide range of plant foods and herbs, come in tens of thousands of different forms, traditionally categorized as flavonoids, tannins, and stilbenes. Major categories of flavonoids include flavonols and flavones (notably quercetin, fisetin, and luteolin), catechins (prominent in green tea and grape seed extract), and anthocyanins (richly supplied by berries and their juices). Tannins, which derive their name from the fact that some have been used in the tanning of leather, are found in pomegranate, raspberries, strawberries, and gall nuts. Stilbenes include the much ballyhooed resveratrol, as well as the up-and-coming pterostilbene, which may have much greater promise for health promotion.

In rodents and cell culture studies, a number of polyphenols have demonstrated positive effects on cognitive function and on the integrity of cultured neurons. These effects include:

- Suppression of microglial inflammation, as reported for anthocyanins from blueberries or acai, the chief catechin in green tea (epigallocatechingallate, or EGCG), fisetin, luteolin, and pterostilbene;²⁷²⁻²⁷⁹
- Improved cognitive function in aging normal mice, seen with green tea, blueberries, and luteolin;^{277, 280-283}
- Improved cognitive function in AD model mice, as demonstrated with green tea catechins, anthocyanins from blueberry or mulberry, grape seed extract, tannic acid, pomegranate juice, and pterostilbene;²⁸⁴⁻²⁹¹
- Protection of neurons from the toxic effects of amyloid beta in cell culture, as seen with quercetin, EGCG, blueberry extract, and the prominent anthocyanin cyanidin-3-glucoside;²⁹²⁻²⁹⁷
- Improved cognitive function in healthy young rodents, as reported for fisetin, epicatechin, and blueberry anthocyanins.²⁹⁸⁻³⁰⁰

With respect to AD, there is evidence that some polyphenols have the potential to suppress production of the dangerous amyloid beta proteins. Amyloid beta is generated by processing of the large membrane protein amyloid precursor protein (APP), produced by most cells, including those in the brain. Amyloid beta arises when APP is successively cleaved by the membrane-associated enzymes beta-secretase and gamma-secretase; the resulting small peptides, 40-42 amino acids in length, have a tendency to bind to each other, forming small complexes known as oligomers. These oligomers of amyloid beta appear to be the truly dangerous actors in AD.³⁰¹ They act on neurons, directly or indirectly, to induce oxidative stress, pro-inflammatory signaling, and neuronal dysfunction. In particular, they disrupt the complex intraneuronal mechanisms required for long term potentiation (LTP), a process crucial to memory formation. In higher concentrations, they can directly kill neurons, and may contribute to the die-off of neurons seen in late-stage AD. Amyloid beta oligomers can also perturb the function of the brain's glial cells, promoting inflammation in microglia and limiting the ability of astrocytes to take up the excitatory neurotransmitter glutamate. This latter effect is important, because excessive extracellular levels of glutamate can lead to a phenomenon known as excitotoxicity which can induce neuronal dysfunction and death. Hence, via their impact on brain glial cells, amyloid beta oligomers can act *indirectly* to threaten the function and survival of neurons. As noted, amyloid beta oligomers also perturb the cerebral vascular system, impairing the delivery of oxygen and nutrients to affected regions of the brain; this lapse of blood flow, in turn, is believed to contribute to the inflammatory syndrome in AD, possibly boosted amyloid beta production. It is thought that, as AD progresses, all of these effects interact to induce a catastrophic

deterioration of brain structure and function. Although amyloid beta oligomers are now believed to be the chief mediators of amyloid beta's pathogenic impact, amyloid beta can also coalesce into large extracellular aggregates known as plaques; the accumulation of these insoluble plaques can be observed via microscopy, and is a hallmark of AD.

Fortunately, the APP protein can have a benign, alternative fate. If cleaved by the membrane enzyme alpha-secretase before beta-secretase can act on it, APP cannot give rise to amyloid beta. The rate at which amyloid beta is produced is therefore dependent on the the balance between alpha-secretase and beta-secretase activity directed toward APP. One of the most intriguing discoveries about the EGCG found in green tea is that, in concentrations which appear to be physiologically feasible, EGCG selectively increases the brain's alpha-secretase activity, such that APP tends to be more preferentially cleaved by alpha-secretase, and that the production of amyloid beta oligomers is suppressed.^{284, 302-306} This effect has been observed both in neuronal cell cultures and in the brains of mice fed EGCG. Conversely, there is recent evidence that tannins, including tannic acid and the punicalagin (an ellagitannin) that is a prominent component of pomegranate juice, can decrease amyloid beta production by direct inhibition of beta secretase activity.^{289, 307, 308} Unfortunately, these studies may be of little clinical relevance, as there is evidence that these compounds are not absorbed intact; the anti-inflammatory properties of ingested pomegranate juice appear to reflect the actions of urolithins, produced by gut bacteria as a metabolite of punicalagin.³⁰⁹ Nonetheless, there is evidence that oral administration of pomegranate juice does indeed suppress amyloid beta generation, inhibit microglial activation, and improve cognitive function in AD model mice³¹⁰ – so urolithins themselves may be importantly protective, even though their direct effects on neurons or microglia haven't yet been studied.

As noted, some flavonoids have shown the ability to boost cognitive function *in healthy young rodents*. Prominent in this regard is fisetin, which, in concentrations which appear to be physiologically relevant, can amplify the sensitivity of the LTP process in neurons from the brain's hippocampus; this likely explains why feeding fisetin to young rats acutely enhances their performance on standard tests of hippocampus-dependent memory formation.^{298, 311, 312} Other studies have found that chronic feeding of epicatechin or blueberry anthocyanins can have a favorable impact on memory function in young rodents.^{299, 300}

A key reason we can be relatively confident that these remarkable findings in rodents and in cell culture studies are pertinent to humans is the growing epidemiology focusing on green tea. The Tsurugaya Project evaluated cognitive function in over 1,000 Japanese subjects aged 70 or above, using a standard test called the Mini-Mental State Examination.³¹³ Using a score under 26 as a cutoff point for cognitive impairment, those who drank at least 2 cups of green tea daily *were less than half a likely to be cognitively impaired* than those who drank no more than 3 cups a week – a finding that had extremely high statistical significance. Those drinking an intermediate amount of green tea showed an intermediate level of protection. Green tea may also provide cognitive protection by lessening risk for stroke; a number studies from Japanese or China find lower risk for stroke in people who are heavy regular users of green tea.³¹⁴⁻³¹⁸ (This protection from stroke may reflect a favorable effect of EGCG on the endothelial lining of the cerebral arteries, as discussed below.) But perhaps the most impressive recent Japanese epidemiology dealing with green tea derives from the Ohsaki Cohort 2006 Study, which enrolled nearly 14,000 Japanese subjects 65 years old or older, and determined their characteristic consumption of green tea at the time of enrollment.³¹⁹ The researchers then followed these subjects for the next 3 years to

determine which of them became functionally disabled during this time, and correlated this information with their green tea drinking habits. (Functional disability was defined objectively as application for Long Term Care Insurance, a government program that provides daily living assistance for the elderly; disability could result from such common causes as stroke, onset of dementia and severe bone fractures.) As compared to those drinking less than one cup of green tea daily, those drinking 1-2 cups daily were 10% less likely, those drinking 3-4 cups daily were 25% less likely, and those drinking 5 or more cups daily were 33% less likely to become functionally disabled during the 3 years of follow-up. It would be hard to make a stronger case for use of green tea by the elderly!

Unfortunately, it is not possible to find comparable epidemiology addressing other types of dietary polyphenols with protective potential, as consumption of blueberry or pomegranate juice, or of polyphenol supplements, has never (or at least not yet!) achieved the fetishistic intensity of green tea consumption. Nonetheless, a handful of epidemiological surveys from Western countries suggest that even the modest levels of food polyphenols provided by ordinary self-selected diets may have a meaningful impact on cognitive function in the elderly. Researchers in France estimated daily dietary intake of total flavonoids in over 1600 volunteers aged 65 years or older; they then examined the cognitive function of these subjects during ten years of follow-up, using the Mini-Mental State Examination.³²⁰ They found that cognitive loss tended to be significantly greater in those whose flavonoid intake was in the bottom fourth of the distribution (average loss of 2.1 points on the test), as compared to those whose flavonoid intake was in the top fourth (average loss of 1.2 points). A more recent U.S study focused on berry consumption (the chief determinant of dietary anthocyanin intake) in over 16,000 participants in the Nurses' Health Study, and correlated this with subsequent performance on standardized tests of cognitive performance.³²¹ With respect to both blueberries and strawberries, relatively high consumption correlated with a slower rate of cognitive decline; indeed, cognitive decline appeared to be decelerated by about 2.5 years in those with high berry intake. Arguably, those who choose to include ample amount of flavonoid-rich foods in their diets may have other dietary and health habits that are partially responsible for the protection observed in these studies; nonetheless, in the light of the favorable impacts of flavonoids on cognitive function in rodents, these findings are provocative and quite possibly meaningful.

Polyphenols with cognitive-protective potential are found in a wide variety of plant foods, and a range of nutraceuticals featuring such polyphenols are also becoming available. Among flavonoid sources, green tea and green tea polyphenol supplements should take pride of place, in light of the rather astounding epidemiology now coming from Japan. A traditional Japanese teacup is said to contain about 80 mg of catechins (primarily EGCG), and optimal protection is seen in those taking 5-6 cups daily, so a daily catechin intake of about 500 mg, preferably dispersed throughout the day (to mimic the typical ingestion pattern of green tea) would seem to be a prudent target. For anthocyanins, frequent consumption of blueberry juice (also available as blueberry juice concentrate, that can be blended with other fluids) is a feasible strategy. Flavonoids can also be obtained from Concord grape juice (which cuts the tartness of blueberry juice a bit if you blend them), and from nutraceuticals featuring fisetin, quercetin, and grape seed extract. For tannins, pomegranate juice is the best practical source, albeit supplements of tannic acid may become available in the future. With respect to the stilbenes, resveratrol unfortunately is metabolized too rapidly by humans to be of much practical benefit, but its close relative pterostilbene can achieve far higher concentrations in the blood, and is now being studied as a nutraceutical in pilot clinical studies.^{322, 323} A smart policy may be to "think cognitive" when consuming fluids – get most of your fluid

throughout the day from coffee, green tea, and grape, berry, or pomegranate juices, and complement this intake with nutraceutical supplements featuring high-potency polyphenols.

Amplify Nitric Oxide Bioactivity – Nitric oxide is a gas produced by one of three forms of an enzyme known as nitric oxide synthase (NOS). The endothelial form of this enzyme, eNOS, is found in the endothelial cells that line the cerebral arteries, and the NO it produces promotes appropriate blood flow to the brain by exerting a vasodilatory effect, while also suppressing arterial inflammation and helping to prevent blood clots; effective eNOS activity appears to be crucial for stroke prevention.²¹ Neurons contain their own form of NOS, neuronal NOS (nNOS), whose activity is crucial for the LTP process required for memory formation.³²⁴ A third form of NOS, so-called inducible NOS (iNOS), is found in inflammatory cells such as activated microglia; as opposed to eNOS and nNOS, which produce low, non-toxic concentrations of NO, iNOS has the potential to produce high concentrations of NO that can react with the oxidant molecular superoxide to generate the highly toxic compound peroxynitrite; this is capable of killing or damaging neurons and may contribute to the neuronal damage seen in Parkinson's disease and AD.³²⁵

Remarkably, there is recent evidence that the NO produced by eNOS in the cerebral vasculature may act directly on the brain to suppress brain production of amyloid beta; in mice that have been genetically altered to lack eNOS activity, their brains express higher levels of both APP and beta-secretase, and they make higher levels of amyloid beta.³²⁶ Conceivably, this helps to rationalize evidence that risk factors for stroke – which often compromise cerebrovascular eNOS activity – also tend to be risk factors for AD.^{21, 326, 327}

The vital biological activities of the low concentrations of NO produced by eNOS and nNOS are mediated largely by activation of an enzyme that produces the compound cyclic GMP (cGMP). cGMP plays a key role in maintaining effective cerebral blood flow, warding off stroke, and supporting effective memory formation via the process of LTP.^{324, 328} cGMP is also produced in inflamed microglia expressing iNOS; this cGMP acts as a negative feedback signal, suppressing some of the pro-inflammatory activities of microglia, while supporting their protective activity as phagocytes.^{329, 330} Evidently, effective cGMP activity is beneficial for brain health in many respects. The concentration of cGMP is regulated by phosphodiesterase enzymes which break it down. One of these enzymes, PDE5, is inhibited by Viagra (sildenafil) and related drugs prescribed for treatment of erectile dysfunction; these drugs function to boost tissue levels of cGMP, a key mediator of erections. Intriguingly, the Chinese medicinal herb epimedium (a.k.a. horny goat weed) contains a compound, icariin, that can also inhibit PDE5, albeit less potently.³³¹⁻³³³ When ingested in feasible amounts, icariin does not have the acute impact on PDE5 that one sees with Viagra or Cialis, but rodent studies suggest that chronic consumption over days can indeed promote erectile function and boost cGMP levels; the data suggest that decreased expression of PDE5 and/or increased expression of nNOS may play a role in this effect. Whether icariin can be effective for boosting cGMP in humans has not yet been established, albeit icariin nutraceuticals are now available, and anecdotal comments suggest that high-dose icariin may indeed be bioactive in humans; evidently, controlled clinical studies are required to assess this.

One of the adverse effects of amyloid beta on neuronal function is to interfere with the production of cGMP – this is one of the ways in which amyloid beta disrupts LTP.³³⁴⁻³³⁶ PDE5 helps to regulate cGMP in regions of the brain vital for cognition, and both sildenafil and icariin, unlike some drugs, have access

to the brain. In light of the foregoing discussion, it isn't surprising that both of these compounds have been reported to aid cognitive function in AD model or senescence-accelerated mice.³³⁷⁻³⁴³ Both sildenafil and icariin can also suppress the activation of microglia in cell cultures.^{344, 345} How these agents influence early cognitive dysfunction in humans remains to be assessed. Those who wish to try icariin should be aware that the highest potency (and hence most credible) icariin supplements currently available provide 180 mg icariin per capsule, and the recommended dose is 3 capsules daily.

As noted, optimal eNOS activity in the cerebral vasculature helps to maintain effective cerebral blood flow and is crucial for stroke prevention.²¹ One key to maintaining this activity is to control oxidative stress in cerebrovascular endothelial cells, as oxidants can inhibit the activity of eNOS – turning it into an enzyme that generates superoxide rather than protective NO – and superoxide can react directly with NO, destroying its activity and producing dangerous peroxynitrite in the process.³⁴⁶ Owing to its ability to inhibit NADPH oxidase, the PhyCB in spirulina is likely to have particular merit for protecting eNOS bioactivity.²¹

But there are also some flavonoids which can act directly on vascular endothelium to stimulate eNOS activity, boosting NO production. These include epicatechin, the key compound in raw cocoa powder which mediates its ability to acutely increase brain blood flow, and thought to be largely responsible for the absence of hypertension and the superior vascular health enjoyed by the Kuna Indians of Panama, whose traditional diet incorporates about 4 cups of raw cocoa daily.³⁴⁷⁻³⁴⁹ Quercetin, the most common flavonol in the American diet, has a direct activating effect on eNOS comparable to that of epicatechin.³⁵⁰ Although the EGCG in green tea does not acutely stimulate eNOS, its consumption somehow makes vascular eNOS function more effective – likely accounting for the diminished stroke risk of green tea drinkers.³⁵¹⁻³⁵³ Quercetin and green tea polyphenols are of course available as nutraceutical supplements, and specially processed cocoa powder that retains its native epicatechin content is also now available commercially. (Contrary to much current hype, most dark chocolate products on the market today are poor sources of epicatechin.)

Histone Deacetylase Inhibitors – Histone deacetylases (HDACs) are enzymes which work in the nucleus of cells to modulate gene expression by regulating transcription of DNA; by removing acetyl groups from a family of DNA-associated proteins known as histones, they tend to repress the synthesis of messenger RNAs required for new protein synthesis. The late phase of LTP required for long-term memory formation is dependent on induced production of a number of proteins that effectively strengthen the synaptic connections between neurons.³⁵⁴ There is exciting recent evidence that the enzyme HDAC2 functions to antagonize the production of a number of these proteins and that, in addition, the adverse effect of amyloid beta and AD on LTP is largely traceable to increased activity of HDAC2.³⁵⁵⁻³⁵⁸

HDAC2 is one of the so-called type 1 HDACs that are susceptible to inhibition by certain drugs or metabolites that are currently available.³⁵⁹ The expensive cancer drug vorinostat has this activity, as do the anti-epileptic drugs valproate, the orphan drug phenylbutyrate (used to treat a rare metabolic disorder), and the natural metabolite butyrate. Recent studies have concluded that type1 HDAC inhibitors can markedly aid memory function in AD model mice, and may even boost memory function in healthy young mice.^{354, 359-364} It is also possible that this strategy might slow the neuronal die-off characteristic of late-stage AD, as one of the proteins whose synthesis is induced by LTP is BDNF, the hormone evoked by exercise training that promotes neuronal survival.^{357, 365}

So far, there have been no clinical trials to determine whether type 1 HDAC inhibitors can improve memory formation in patients with early-stage cognitive dysfunction. And there are logistical difficulties with current HDAC inhibitors. Vorinostat and phenylbutyrate are extremely expensive – too expensive for off-label use by people who aren't wealthy - and valproate induces tiredness. Butyrate, a safe natural metabolite, has a blood half life of only about 6 minutes, as it is rapidly burned as fuel by the body's tissues; hence, unless ingested constantly, it presumably would be of limited value. A ray of hope is offered by the fact that tributyrin (glyceryl-tributyrate) is an approved food additive that is broken down gradually in the body to act as a time-release source of butyrate.³⁶⁶⁻³⁶⁹ A decade or so ago, tributyrin was tested in cancer patients, as HDAC inhibitors have some potential in cancer therapy (witness the recently approved drug vorinostat).³⁶⁷ But tributyrin would need to be administered in very high daily doses to be effective – perhaps several tablespoons daily - and in these clinical studies it was administered in capsules, entailing the ingestion of 60 or more capsules daily! This obviously would be a non-starter from a compliance standpoint. Hence, tributyrin will not be clinically practical until and unless nutraceutical entrepreneurs find a feasible way to incorporate high doses in functional foods.

In any case, let's hope that phenylbutyrate is soon evaluated in patients with early-stage cognitive dysfunction sometime, so we can learn whether this avenue of therapy is worth pursuing; this agent is currently being tested clinically in several other neurodegenerative disorders, and is well tolerated in doses of 9-15 grams daily.^{370, 371}

One of carnitine's physiological roles is to shuttle acetyl groups from mitochondria to the nucleus, which promotes the acetylation of histones and other nuclear proteins.³⁷²⁻³⁷⁵ Evidently, this has the potential to counteract the adverse impact of HDAC2 activity on memory formation. So it is reasonable to suspect that some of acetyl-L-carnitine's favorable impact in early Alzheimer's reflects opposition to HDAC2. Concurrent administration of ALC may therefore boost any favorable impact of histone deacetylase inhibitors such as tributyrin on memory function in early AD.

Targeting Tau - Thus far, little has been said in this essay about tau protein, whose excessive phosphorylation in AD and certain other dementing neurodegenerative conditions disrupts its function and leads to the formation of abnormal intraneuronal structure known as neurofibrillar tangles. Amyloid beta drives the hyperphosphorylation of tau, and this phosphorylation of tau is believed to play a key mediating role in AD neurodegeneration; indeed, AD model mice which are also bioengineered to produce lesser amounts of tau are protected from cognitive decline.³⁷⁶⁻³⁷⁸ Moreover, hyperphosphorylation of tau plays a pathogenic role in other types of dementing neurodegeneration not linked to amyloid beta excess, such as frontotemporal dementia (these are known as "tauopathies"). Hence, measures which can prevent or reverse the phosphorylation of tau are being sought as therapeutic options.

One of the key enzymes which phosphorylates tau in AD is glycogen kinase synthase-3beta (GSK-3 β). It has long been suspected that the therapeutic efficacy of lithium in manic-depressive illness hinges on its ability to inhibit this enzyme;³⁷⁹ it is therefore reasonable to suspect that therapeutic doses of lithium might lessen tau phosphorylation in AD or other neurodegenerative conditions. Moreover, lithium may also have the potential to decrease amyloid beta production, since phosphorylation of APP by GSK-3 β increases its propensity to be converted to amyloid beta.^{380, 381} Indeed, several epidemiological studies have found that, in patients with manic-depressive disorder, those receiving lithium are at much lower

risk for AD than those not receiving this drug.³⁸²⁻³⁸⁴ This has been followed up with a controlled 12-month pilot study in which, while both groups experienced further cognitive decline, this was of lesser magnitude in the lithium-treated group.³⁸⁵ Lithium has also shown a favorable effect on cognitive function in some, but not all,^{386, 387} studies in AD model mice, and has reduced the adverse impact of amyloid beta in neuronal cell cultures.^{380, 388, 389} And a very recent report suggests that “mini-doses” of lithium (300 mcg daily) may also be useful in early cognitive decline;³⁹⁰ standard dosing with lithium is potentially toxic and must be carefully monitored by a physician to insure that blood levels do not exceed the safe range.

An enzyme which de-phosphorylates tau (and hence potentially mitigates the problem) is protein phosphatase 2A (PP2A). Remarkably, it has recently been discovered that the compound sodium selenate – long employed as a supplemental source of selenium – can interact with PP2A in a way that boosts its ability to dephosphorylate tau.^{391, 392} In a strain of mouse bioengineered to overexpress a mutant human form of tau that gives rise to neurofibrillary tangles, simply giving selenate in drinking water (12 mg per liter) was associated with a marked reduction in tau phosphorylation, a complete prevention of neurofibrillary tangles, and an improvement in memory. While the mice appeared to tolerate this treatment well, selenium is potentially toxic, and, though selenate is the best tolerated form of supplemental selenium, it remains to be established that prolonged administration of selenate doses sufficient to reduce tau phosphorylation can be tolerated well by humans. Fortunately, a recent phase I clinical study, in which sodium selenate supplementation was assessed in prostate cancer patients, has reported a maximum tolerated daily dose of 60 mg, which is likely more than would be required to modulate PP2A activity.³⁹³ Hence, selenate may ultimately emerge as a useful option in the management of AD and other neurodegenerative conditions associated with tau hyperphosphorylation. And lithium plus selenate might supply a “one-two punch” in that regard. Arguably, such a strategy would have the best chance for important clinical benefit if allied with additional measures that more directly address amyloid beta toxicity.³⁸⁶

Inhibiting 5-Lipoxygenase – The omega-6 fatty acid arachidonic acid can be enzymatically converted to a number of pro-inflammatory factors, known as prostanoids and leukotrienes. The cyclooxygenase enzymes give rise to the prostanoids, and NSAID drugs commonly used to treat inflammation – agents such as aspirin, ibuprofen, and Celebrex – are potent inhibitors of these enzymes. Some epidemiological studies have found that people who have been chronic users of NSAID drugs (people with rheumatoid arthritis, for example) are at markedly lower subsequent risk for AD.³⁹⁴⁻³⁹⁶ This encouraged clinical trials of NSAIDs in patients suffering the early stages of AD; unfortunately, no benefit for cognitive function was seen, and the patients receiving the NSAIDs were at higher risk for side effects and cardiovascular mortality.³⁹⁷ It seems that brain prostanoids can help to trigger an inflammatory process that, once florid, has its own self-sustaining momentum. NSAIDs may therefore have some utility for the prevention of AD, but cannot be recommended for its treatment. And unfortunately, the side effects and increased cardiovascular risk associated with NSAID therapy (especially with the putatively less toxic cyclooxygenase-2-specific inhibitors) makes it imprudent to recommend these agents for AD prevention.

However, there is now evidence which suggests that another pro-inflammatory enzyme, 5-lipoxygenase, may also contribute to the progression of AD. 5-lipoxygenase (5-LOX) produces pro-inflammatory factors known as leukotrienes. 5-LOX expression has been found to be increased in the brain (specifically in the hippocampus, the key target of AD inflammation) of people with AD, possibly owing

to induction by stress hormones.^{398, 399} In mouse models of AD, measures which increase the expression of 5-LOX within the brain exacerbate memory deficits, boost the production of amyloid beta, and increase the phosphorylation of tau.^{399, 400} This increased production of amyloid beta has been traced to increased brain expression of gamma-secretase, which plays an essential catalytic role in this production; and increased tau phosphorylation reflects increased activity of the enzyme CDK5, whose activation in neurons is a common feature of AD. In cell culture studies, increased 5-LOX activity makes neuron-like cells more vulnerable to the toxic impact of amyloid beta.⁴⁰¹ Conversely, drugs which inhibit 5-LOX activity, or genetic measures which prevent 5-LOX expression in the brain, tend to ameliorate the memory decline, suppress amyloid beta production, and lessen tau phosphorylation in AD model mice.⁴⁰²⁻⁴⁰⁶

Since 5-LOX plays a key role in asthma, drugs which inhibit it – notably zileuton – are currently available, and, in notable contrast to cyclooxygenase inhibitors, seem to be safe and relatively well tolerated.⁴⁰⁷ Natural compounds known as boswellic acids – richly supplied by a traditional Ayurvedic herbal remedy, salai guggal – also can act as 5-LOX inhibitors, and boswellic acid-enriched extracts are now available as nutraceuticals.⁴⁰⁸⁻⁴¹⁰ However, there is now considerable doubt that the clinical anti-inflammatory activity of boswellic acids reflects 5-LOX inhibition; clinically, it may function to decrease the production of prostaglandin E1, a compound which mediates much of the pro-inflammatory impact of cyclooxygenase.^{411, 412} So perhaps, in some adequate intake, boswellic acids might provide some of the protective benefits of cyclooxygenase inhibitors, without the toxic risk. Joint administration of a form of boswellic acid and a cyclooxygenase-2 inhibitory drug for 15 days was reported to reverse age-related cognitive impairments in mice.⁴¹³

Since the evidence relating AD to 5-LOX activity is very recent, there aren't yet any clinical trials that have assessed the impact of zileuton on early AD; such studies now appear to be warranted. It has been suggested that a possible reason why NSAIDs haven't proven useful in treating AD is that these drugs tend to make arachidonic acid more available to 5-LOX, potentiating production of leukotrienes.⁴¹⁴ Intriguingly It should be noted that diets with a very high ratio of omega-3 to omega-6 can suppress production of pro-inflammatory leukotrienes by lessening arachidonic acid availability.⁴¹⁵

Supporting Synaptogenesis – As we noted above, one of the key roles of DHA is to support the efficient production of new synaptic membranes required for memory formation. But DHA is not the only nutrient that can be rate-limiting in this regard. Richard Wurtman and colleagues, working with gerbils, have shown that brain availability of cytidine – a key cofactor required for the production of the phospholipids that constitute the bulk of neuronal membranes – can also be rate-limiting for synaptogenesis. Although the brain does not take up cytidine efficiently from this blood, it does efficiently take up uridine, which the brain rapidly converts to cytidine.⁴¹⁶ Wurtman's studies show that supplementing gerbils with uridine (as uridine monophosphate, an approved food additive) improves their performance in memory tests; this benefit is potentiated if DHA is administered concurrently.^{99, 417, 418} Moreover, this strategy is associated with an increased level of dendritic spines – the receptive component of a synapse - in the hippocampus of these animals. Bioavailable sources of choline (see below) can also be beneficial in this regard, as choline is also a precursor for phospholipid synthesis. Hence, Wurtman recommends DHA, uridine monophosphate, and bioavailable choline as a strategy for improving memory function; clinical studies to evaluate this are underway.⁴¹⁹

The compound orotate is of related interest. Orotate is a natural metabolite that, when ingested orally, is rapidly taken up by the liver and converted primarily to uridine, most of which is then returned to the bloodstream.^{420, 421} Hence, orotate might function as an alternative to uridine monophosphate. Intriguingly, in a number of studies dating back to the 1970s and 1980s, European researchers reported that various complexes of orotate had favorable effects on memory in rodent models of cognitive dysfunction.⁴²²⁻⁴²⁵ Magnesium orotate is of particular interest, as it appears to be clinically useful in congestive heart failure and in angina, common conditions in the elderly; uridine is likely to be the primary mediator of this benefit.⁴²⁶⁻⁴²⁸ The dose of magnesium orotate which appears to benefit congestive failure is 3-6 g daily; if this dose produces enough uridine to benefit the heart, it likely also would produce enough to influence brain function.

Increasing Brain Magnesium - Intriguingly, there are recent reports that supplementation which boosts brain levels of magnesium may also aid synaptogenesis. The compound threonate (a metabolite of vitamin C) may aid transport of magnesium across the blood brain barrier and increase the concentration of magnesium in cerebrospinal fluid.^{429, 430} This effect has been associated with increased synaptogenesis and improved learning in rats. More recently, magnesium threonate supplementation throughout life has been reported to have a markedly favorable impact on AD model mice – reducing brain levels of soluble amyloid beta, preventing alterations in key brain signaling molecules, prolonging their survival, and most importantly normalizing their performance in certain cognitive tests.⁴³¹ And in older AD model mice, one month of magnesium threonate had a very positive impact on their cognitive performance. Another report indicates that, in neuronal cell cultures, a slightly supraphysiological concentration of magnesium promotes the activity of alpha-secretase, and hence impedes amyloid beta generation.⁴³² Other researchers have shown that increased extracellular levels of magnesium can dampen the activation of microglia, whose excessive activation may play a role in AD, Parkinson's, and other neurodegenerative conditions.⁴³³ And there are several reports that brain and plasma magnesium levels tend to be low in patients with AD.⁴³⁴⁻⁴³⁶ These findings all point to the possibility that measures which increase brain magnesium level may be importantly neuroprotective - possibly because they help to prevent excessive pro-inflammatory uptake of calcium by neurons and microglia.

Unfortunately, there are not yet any published reports assessing the impact of magnesium threonate in humans; however, this compound is now commercially available as a supplement, and likely is quite safe. Whether threonate is unique in its ability to aid transport of magnesium into the brain remains to be seen; orotate, which has traditionally been viewed as a “mineral transporter”, has not been evaluated in this regard.

Boosting Cholinergic Activity – The progression of AD tends to selectively kill and damage “cholinergic” neurons which function to release the neurotransmitter acetylcholine; loss of acetylcholine activity is responsible for much of the cognitive dysfunction in AD. Two compensatory strategies are feasible – providing more choline to the brain, so that the remaining viable cholinergic neurons can make acetylcholine more efficiently, and inhibiting the acetylcholinesterase enzyme that terminates acetylcholine's bioactivity by degrading it. Each of these strategies has been avidly pursued clinically.

Taking straight choline is not a viable way increase brain choline levels, as free choline is inefficiently absorbed, and the unabsorbed choline is degraded by gut bacteria to generate a “dead fish” odor. The nutraceutical delivery vehicles for choline that have shown the most promise are

glycerylphosphorylcholine (a.k.a. GPC or choline alfoscerate) and CDP-choline (a.k.a. citicoline).⁴³⁷ The former has received more clinical evaluation from the standpoint of cognitive impairment, whereas citicoline has been studied primarily from the standpoint of its ability to restore brain phospholipid levels following a stroke.⁴³⁸ An overview of clinical trials with GPC suggests that it is indeed modestly beneficial for improving cognitive function in early-stage cognitive impairment, in a dose of 1200 mg daily.⁴³⁹

While choline is obviously required for acetylcholine synthesis, so is an acetyl group. Acetyl groups are generated in profusion by mitochondrial metabolism, but these need to be shuttled to the cell cytoplasm if they are to be used for acetylcholine synthesis. Carnitine is the carrier for this shuttle system, and this likely explains why acetyl-L-carnitine has been shown to boost the efficiency of acetylcholine synthesis in neurons.²⁵⁴⁻²⁵⁶ This effect no doubt contributes to the improvement of cognitive function achieved with supplemental acetyl-L-carnitine in elderly people with early AD.

The first drug approved for treating AD, donepezil (a.k.a. Aricept), is an acetylcholinesterase inhibitor; other agents of this type that have been approved are tacrine, rivastigmine, and galantamine. Subsequently, scientists discovered that huperzine A, the active component of the herb *Huperzia serrata* long used to treat age-related cognitive dysfunction in China, likewise can inhibit acetylcholinesterase, and that its pharmacological properties are superior to those of donepezil.⁴⁴⁰ An overview of numerous controlled Chinese clinical trials with huperzine A concludes that, in doses averaging 200 mcg twice daily, it is indeed effective for improving diminished cognitive function, and appears to be at least as effective as donepezil in this regard.⁴⁴¹ Furthermore, studies in neuronal cell cultures and in AD model mice suggest that huperzine A may have some neuroprotective properties not related to its abilities to inhibit acetylcholinesterase.⁴⁴²⁻⁴⁴⁴ Huperzine A is now available as an affordable nutraceutical in the U.S. Acetylcholinesterase inhibitors are usually well tolerated in recommended doses, albeit an associated increase in the cholinergic activity of the GI tract occasionally induces symptoms such as nausea and vomiting.

The downside of choline-boosting strategies for cognitive dysfunction is that they do not affect the implacable underlying inflammatory process associated with AD, and they gradually lose their efficacy as the die-off of the cholinergic neurons becomes so extensive that the remaining viable neurons can generate too little acetylcholine to be worthwhile. Also, there does not appear to be any evidence that selective loss of cholinergic neurons is part of the normal aging process, so these agents may be primarily useful in early AD.⁴⁴⁵ However, the possibility that huperzine A may confer neuroprotection independent of its impact on acetylcholine merits further evaluation.

Decreasing Excitotoxicity – As noted above, the chronic stimulation of certain neuronal receptors (extrasynaptic NMDA receptors) by inappropriately elevated extracellular levels of the neurotransmitter glutamate is now suspected to be a major way in which amyloid beta perturbs the function and jeopardizes the survival of neurons; this phenomenon is known as excitotoxicity and, in a starker and more acute way, also contributes to the brain damage inflicted by stroke. Caffeine may be beneficial in this regard by aiding control of extracellular glutamate.⁴⁴⁶ However, the drug memantine, via suppressive interaction with NMDA receptors, can also aid in control of excitotoxicity, and has been demonstrated to slow cognitive decline in patients with mid-stage AD.⁴⁴⁷ Indeed, aside from a group of cholinesterase inhibitor drugs of which donepezil is the prototype, memantine is the only other drug currently approved

for use in AD treatment. Although it is only modestly beneficial for cognition, it does appear to slow subsequent atrophy of the hippocampus, and hence may be a quite worthwhile option as a follow up to cholinesterase treatment.^{448, 449} It does not seem to be beneficial in early-stage AD, possibly because memantine only blocks NMDA receptors when they are very strongly activated.^{450, 451} (Indeed, activation of synaptic NMDA receptors is crucial for LTP and memory formation, so memantine would be expected to worsen cognition if its inhibitory activity were non-selective.)

A key mediator of intracellular signaling by extracellular NMDA receptors is activation of the proteolytic enzyme calpain.^{96, 452} Recently a drug inhibitor of calpain has shown remarkable benefit in AD mouse models, without overt toxicity.⁴⁵³⁻⁴⁵⁵ Perhaps this, or some analogous drug, will become available for clinical use in the future.

Neuronal Growth Factor Mimics – A particularly interesting recent development is the discovery by Emory University scientists that the flavonoid 7,8-dihydroxyflavone – which is said to occur naturally in tiny amounts – can directly activate a key neuronal receptor for BDNF, TrkB.⁴⁵⁶ Activation of this receptor plays a key role in support of LTP, and also aids neuronal survival; thus, 7,8-dihydroxyflavone might be expected to have potential for promoting effective cognitive function and staving off neurodegeneration.⁴⁵⁷ Another group soon showed that administration of this molecule reversed memory failure and suppressed expression of beta-secretase and of amyloid beta in an AD mouse model; and this molecule likewise is beneficial for normal age-related cognitive decline in rats.^{458, 459} The Emory group has also identified another natural molecule, deoxygedunin from the neem tree, that can directly activate the TrkB receptor.⁴⁶⁰ These discoveries are particularly exciting in light of the fact that direct administration of the BDNF neural growth factor is impractical, in that it does not penetrate the brain after intravenous administration, and also has a limited half-life.

Meanwhile, other researchers have discovered that sominone, a natural metabolite of withanoside IV found in the ayurvedic herb ashwagandha, can directly activate another neural growth factor receptor, RET.⁴⁶¹ 60 minutes after sominone was injected into normal mice, their memory performance was improved. These researchers soon followed up with evidence that sominone improved memory in AD model mice.⁴⁶² Unfortunately, withanoside IV per se is not currently available as a supplement, and the amount of ashwagandha extract required to provide an effective dose of this compound would likely be quite high.

Another feasible way to increase neuron-supportive growth factor activity in the brain is via intranasal administration of insulin.⁴⁶³ In the many neurons which express insulin receptors, increased insulin activity could be expected to decrease phosphorylation of tau, and also increase the survival of stressed neurons.⁴⁶⁴ Two recent pilot clinical trials have evaluated intranasal insulin (20 to 40 IU daily, for 3 weeks or 4 months) in patients with mild cognitive impairment or early AD; both doses were found to aid preservation of cognitive function and functional ability.^{465, 466} These doses of insulin were well tolerated; they did not detectably influence blood insulin levels and hence caused no side effects related to low blood sugar. Further studies evaluating this intriguingly simple strategy are in progress.

Clearing Amyloid Beta from the Brain - Ashwagandha figures in another provocative recent study. Although much attention has been devoted to strategies to suppress amyloid beta synthesis, there is evidence that the accumulation of amyloid beta in the brains of AD patients may be more attributable to inefficient disposal of amyloid beta than to increased synthesis.⁴⁶⁷ The brain has mechanisms both for

degrading amyloid beta, and for expelling it into the blood stream. One of these latter mechanisms is dependent on expression of the protein LRP1, which is also expressed in the liver and enables uptake and degradation of amyloid beta from the blood.^{468, 469} In a recent study, Indian researchers discovered that massive oral doses of a withanolide-rich ashwagandha extract markedly induced LRP1 levels in the liver of AD model mice, such that blood as well as brain levels of amyloid beta markedly declined.⁴⁷⁰ This effect was associated with behavioral improvements in the mice. Sadly, the amount of ashwagandha extract employed in this study would be rather ruinously expensive if translated into a human dose, so it is not clear that this discovery has any practical import. Nonetheless, it does suggest that a more practical means of inducing LRP1 might help prevent and possibly control AD. A recent report suggested that the drug rifampin could induce increased LRP1 expression in cerebral blood vessels⁴⁷¹ – in nice concordance with a previous clinical study suggesting that it might be useful in early AD⁴⁷² - but unfortunately this drug failed to improve cognitive function in AD patients in a more substantial recent controlled trial.⁴⁷³

The uptake and degradation of amyloid beta fibrils by microglia, which contribute importantly to amyloid beta disposal, is dependent on the production of the apoE protein by astroglia.⁴⁷⁴⁻⁴⁷⁶ Impairment of this mechanism may explain why people who inherit the variant apoE4 form of this protein are at greatly increased risk for AD.⁴⁷⁷ Drugs which activate the LXR receptor, being evaluated as potential treatments for atherosclerosis, promote apoE production in the brain, and in AD mice decrease brain amyloid beta levels while improving cognitive function; these drugs are not currently approved for clinical use, however.^{475, 477-480} Fortuitously, there is a report that the antioxidant nutrient taurine has the potential to at least mildly activate a form of the LXR receptor expressed by astrocytes; it therefore potentially might influence brain amyloid beta metabolism.⁴⁸¹ Taurine has not yet been tested in AD model mice, so this possibility remains hypothetical. The LXR receptor partners with the RXR receptor in inducing apoE, and a cancer drug which activates the RXR receptor, bexarotene, has also been reported to boost apoE expression and reduce amyloid beta levels in AD model mice.⁴⁸² Moreover, the diabetes drugs Actos (pioglitazone) and Avandia (rosiglitazone), via activation of the PPARgamma receptor, can increase expression of the LXR receptor, and hence may act indirectly to increase production of apoE.⁴⁸³ These drugs have shown favorable effects on cognitive function and amyloid beta levels in mouse AD models, and hence may have some potential for AD control.⁴⁸³⁻⁴⁸⁷ Pilot clinical trials evaluating their impact in patients with cognitive impairment have so far yielded mixed results.⁴⁸⁸ In any case, this ongoing research bears watching. In light of evidence that rosiglitazone can increase heart attack risk in diabetics, pioglitazone would likely be the superior choice.^{489, 490}

Overview and Provisional Recommendations

Every now and then we hear about the clinical results with another new drug-in-development intended to treat AD; more often than not, the results are equivocal if not downright disappointing.^{491, 492} The fact is that AD is an extremely complex and still rather poorly understood disorder, for which mouse models are a very imperfect replica, and it may be quite unrealistic to expect any single drug, new or old, to offer definitive and lasting benefit. More likely, effective prevention and treatment of this disorder will require a complex regimen of drug, nutraceutical, and lifestyle measures that address in a complementary manner a range of the dysfunctions that promote and sustain it. And it is increasingly recognized that attempting to control AD when it is already floridly symptomatic may be a losing proposition – AD is likely to be a disorder that it more readily prevented (or at least markedly postponed) than cured.⁴⁹³

Proactive measures are also required for stroke prevention, and for optimal prevention of age-related cognitive dysfunction.

The foregoing discussion provides evidence that quite a number of measures, many of them reasonably practical, have credible potential for warding off cognitive decline and AD during aging. But it clearly would be impractical and quite likely inadvisable to implement all of these measures simultaneously – a triage strategy is evidently needed. I suggest that we classify these measures as first-line – recommended for use by healthy people before symptoms of cognitive dysfunction manifest; second-line – relatively practical measures which can be introduced when signs of cognitive decline first appear (or, in the future, when a doctor can detect incipient AD by novel diagnostic techniques); and third-line – measures which are less inherently practical or affordable, but which can be considered as options if cognitive dysfunction continues to worsen. First-line measures are intended to slow onset of age-related cognitive decline and AD, while decreasing stroke risk; they should also be expected to have a favorable impact on overall health. Provisionally (as no one truly has “the answer” to this vexing health conundrum), I suggest the following:

First-Line Measures – for primary prevention

- Mediterranean or plant-based diet, low in saturated fat, high in fruits and vegetables, moderate in salt and high in potassium;
- Aerobic exercise training
- Avoidance of obesity and diabetes (which should be aided greatly by the previous two measures);
- Anti-hypertensive therapy as needed (brain-permeable angiotensin antagonists preferred);
- Effective antioxidant supplementation, including: spirulina – 15 g/day; astaxanthin – 4-20 mg/d; lipoic acid – 600-1200 mg/d; NAC or cystine – 600-1800 mg/d; taurine – 2-4 g/d
- Caffeinated coffee – as much and as often as you can without impairing effective sleep or inducing other unacceptable side effects
- Green tea and green tea polyphenols – several cups daily and/or several caps of green tea polyphenol extract (at least 500 mg of green tea catechins daily)
- Blueberry, pomegranate, and Concord grape juices, multiple servings daily (you may blend them if desired);
- DHA, 500-1,000 mg daily (from supplements or frequent ingestion of oily fish)
- Quercetin or cocoa flavanols, several times daily
- Magnesium threonate, up to 400 mg magnesium daily.

Second-Line Measures – To be added to first-line measures when cognitive dysfunction first appears

- Acetyl-L-Carnitine – 2-3 g daily
- Melatonin – 3-10 mg at bedtime
- Fisetin – 200-300 mg twice daily
- Icaritin – 180 mg, 3 times daily
- Magnesium Orotate – 2-4 g daily
- Glycerylphosphorylcholine (GPC) 400 mg – 3 caps daily; and/or Huperzine A – 200 mcg twice daily
- Zileuton – 1200 mg, twice daily

- Inosine – 1-3 g daily (for incipient PD or DLB – dose to be titrated with physician supervision to insure that serum urate does not exceed 9 mg/dL; alkalinizing diet advisable)

Third-Line Measures – Options to consider as cognitive dysfunction continues to progress, with the cooperation and supervision of your physician

- Memantine – gradually tritrated up to 20 mg daily
- Lithium – dosage intended to maintain blood level of 0.25- 0.5 mM, as determined by your doctor (or recently described 300 mcg/d mini-dose regimen)
- Sodium selenate – 20-30 mg daily (note – potential for side effects)
- Tributyrin – 10 g 3 times daily (when and if available)
- Pioglitazone – 30 mg (may cause fluid retention; contraindicated in congestive heart failure)
- Intranasal Insulin – 20 IU once daily

This list is not exhaustive, and is admittedly somewhat subjective – no doubt other competent analysts would provide recommendations that differed in some particulars. The roster of phytochemicals which have been assessed in rodent models of cognition is very long and getting longer; it wouldn't be logistically feasible to take meaningful doses of all of the phytochemicals which have shown promise in these studies. (Those who would like to add grape seed extract, ginseng, ginkgo, ashwagandha, bacopa, etc. to their neuroprotective regimens are welcome to do so.⁴⁹⁴⁻⁴⁹⁷) It also should be acknowledged that many of the recommended measures might not be notably effective if tested as monotherapies in AD; rather, the hope is that, by addressing a number of different aspects of the AD syndrome simultaneously, the combination of these measures will provide worthwhile benefit. **In aggregate, these measures are intended to dampen microglial activation, provide comprehensive antioxidant support for healthful brain function, boost protective NO/cGMP function, block A2A adenosine receptors, inhibit leukotriene production, suppress amyloid beta synthesis, promote microglial disposal of amyloid beta, inhibit HDAC2 activity, inhibit tau phosphorylation, suppress the production and pathogenic impact of peroxynitrite, potentiate LTP, support acetylcholine activity, support efficient synaptogenesis, increase brain magnesium levels, decrease excitotoxicity, promote efficient cerebral blood flow, and lessen stroke risk.**

It is greatly to be hoped that within the next few years one or more promising new drugs (or possibly 7,8-dihydroxyflavone?) will become available for AD treatment. But it would be rash to sit back complacently and expect that some future new drug will be “the answer” to AD. The smart thing to do is to start taking steps to protect your aging brain when you are still relatively young – or at least before frank cognitive dysfunction manifests. It bears re-emphasis that AD is a disease better prevented than treated. And that's obviously true of stroke as well.

References

- (1) Kalmijn S, Launer LJ, Ott A, Witteman JC, Hofman A, Breteler MM. Dietary fat intake and the risk of incident dementia in the Rotterdam Study. *Ann Neurol* 1997 November;42(5):776-82.
- (2) Gu Y, Luchsinger JA, Stern Y, Scarmeas N. Mediterranean diet, inflammatory and metabolic biomarkers, and risk of Alzheimer's disease. *J Alzheimers Dis* 2010;22(2):483-92.
- (3) Gu Y, Nieves JW, Stern Y, Luchsinger JA, Scarmeas N. Food combination and Alzheimer disease risk: a protective diet. *Arch Neurol* 2010 June;67(6):699-706.
- (4) Morris MC, Evans DA, Bienias JL et al. Dietary fats and the risk of incident Alzheimer disease. *Arch Neurol* 2003 February;60(2):194-200.
- (5) Laitinen MH, Ngandu T, Rovio S et al. Fat intake at midlife and risk of dementia and Alzheimer's disease: a population-based study. *Dement Geriatr Cogn Disord* 2006;22(1):99-107.
- (6) Patil S, Sheng L, Masserang A, Chan C. Palmitic acid-treated astrocytes induce BACE1 upregulation and accumulation of C-terminal fragment of APP in primary cortical neurons. *Neurosci Lett* 2006 October 2;406(1-2):55-9.
- (7) Patil S, Melrose J, Chan C. Involvement of astroglial ceramide in palmitic acid-induced Alzheimer-like changes in primary neurons. *Eur J Neurosci* 2007 October;26(8):2131-41.
- (8) Patil S, Balu D, Melrose J, Chan C. Brain region-specificity of palmitic acid-induced abnormalities associated with Alzheimer's disease. *BMC Res Notes* 2008;1:20.
- (9) Liu L, Martin R, Chan C. Palmitate-activated astrocytes via serine palmitoyltransferase increase BACE1 in primary neurons by sphingomyelinases. *Neurobiol Aging* 2012 June 22.
- (10) Gupta S, Knight AG, Gupta S, Keller JN, Bruce-Keller AJ. Saturated long-chain fatty acids activate inflammatory signaling in astrocytes. *J Neurochem* 2012 March;120(6):1060-71.
- (11) Wang Z, Liu D, Wang F et al. Saturated fatty acids activate microglia via Toll-like receptor 4/NF-kappaB signalling. *Br J Nutr* 2012 January;107(2):229-41.
- (12) Milanski M, Degasperi G, Coope A et al. Saturated fatty acids produce an inflammatory response predominantly through the activation of TLR4 signaling in hypothalamus: implications for the pathogenesis of obesity. *J Neurosci* 2009 January 14;29(2):359-70.
- (13) Ross FM, Allan SM, Rothwell NJ, Verkhratsky A. A dual role for interleukin-1 in LTP in mouse hippocampal slices. *J Neuroimmunol* 2003 November;144(1-2):61-7.
- (14) Vereker E, O'Donnell E, Lynch MA. The inhibitory effect of interleukin-1beta on long-term potentiation is coupled with increased activity of stress-activated protein kinases. *J Neurosci* 2000 September 15;20(18):6811-9.
- (15) Griffin WS, Barger SW. Neuroinflammatory Cytokines-The Common Thread in Alzheimer's Pathogenesis. *US Neurol* 2010;6(2):19-27.

- (16) Krause DL, Muller N. Neuroinflammation, microglia and implications for anti-inflammatory treatment in Alzheimer's disease. *Int J Alzheimers Dis* 2010;2010.
- (17) Rogers J, Mastroeni D, Leonard B, Joyce J, Grover A. Neuroinflammation in Alzheimer's disease and Parkinson's disease: are microglia pathogenic in either disorder? *Int Rev Neurobiol* 2007;82:235-46.
- (18) Griffin R, Nally R, Nolan Y, McCartney Y, Linden J, Lynch MA. The age-related attenuation in long-term potentiation is associated with microglial activation. *J Neurochem* 2006 November;99(4):1263-72.
- (19) Barrientos RM, Frank MG, Watkins LR, Maier SF. Memory impairments in healthy aging: Role of aging-induced microglial sensitization. *Aging Dis* 2010 January 1;1(3):212-31.
- (20) Lindeberg S, Lundh B. Apparent absence of stroke and ischaemic heart disease in a traditional Melanesian island: a clinical study in Kitava. *J Intern Med* 1993 March;233(3):269-75.
- (21) McCarty MF. Up-regulation of endothelial nitric oxide activity as a central strategy for prevention of ischemic stroke - just say NO to stroke! *Med Hypotheses* 2000 November;55(5):386-403.
- (22) Fiocco AJ, Shatenstein B, Ferland G et al. Sodium intake and physical activity impact cognitive maintenance in older adults: the NuAge Study. *Neurobiol Aging* 2012 April;33(4):829-8.
- (23) Khachaturian AS, Zandi PP, Lyketsos CG et al. Antihypertensive medication use and incident Alzheimer disease: the Cache County Study. *Arch Neurol* 2006 May;63(5):686-92.
- (24) Yasar S, Lin FM, Fried LP et al. Diuretic use is associated with better learning and memory in older adults in the Ginkgo Evaluation of Memory Study. *Alzheimers Dement* 2012 May;8(3):188-95.
- (25) Cappuccio FP, MacGregor GA. Does potassium supplementation lower blood pressure? A meta-analysis of published trials. *J Hypertens* 1991 May;9(5):465-73.
- (26) Khaw KT, Barrett-Connor E. Dietary potassium and stroke-associated mortality. A 12-year prospective population study. *N Engl J Med* 1987 January 29;316(5):235-40.
- (27) Bazzano LA, He J, Ogden LG et al. Dietary potassium intake and risk of stroke in US men and women: National Health and Nutrition Examination Survey I epidemiologic follow-up study. *Stroke* 2001 July;32(7):1473-80.
- (28) Iso H, Stampfer MJ, Manson JE et al. Prospective study of calcium, potassium, and magnesium intake and risk of stroke in women. *Stroke* 1999 September;30(9):1772-9.
- (29) He FJ, MacGregor GA. Potassium: more beneficial effects. *Climacteric* 2003 October;6 Suppl 3:36-48.
- (30) Green DM, Ropper AH, Kronmal RA, Psaty BM, Burke GL. Serum potassium level and dietary potassium intake as risk factors for stroke. *Neurology* 2002 August 13;59(3):314-20.

- (31) Larsson SC, Virtamo J, Wolk A. Potassium, calcium, and magnesium intakes and risk of stroke in women. *Am J Epidemiol* 2011 July 1;174(1):35-43.
- (32) O'Donnell MJ, Yusuf S, Mentz A et al. Urinary sodium and potassium excretion and risk of cardiovascular events. *JAMA* 2011 November 23;306(20):2229-38.
- (33) D'Elia L, Barba G, Cappuccio FP, Strazzullo P. Potassium intake, stroke, and cardiovascular disease a meta-analysis of prospective studies. *J Am Coll Cardiol* 2011 March 8;57(10):1210-9.
- (34) McCabe RD, Bakarich MA, Srivastava K, Young DB. Potassium inhibits free radical formation. *Hypertension* 1994 July;24(1):77-82.
- (35) Ishimitsu T, Tobian L, Sugimoto K, Everson T. High potassium diets reduce vascular and plasma lipid peroxides in stroke-prone spontaneously hypertensive rats. *Clin Exp Hypertens* 1996 July;18(5):659-73.
- (36) McCarty MF. Endothelial membrane potential regulates production of both nitric oxide and superoxide--a fundamental determinant of vascular health. *Med Hypotheses* 1999 October;53(4):277-89.
- (37) Umesawa M, Iso H, Date C et al. Relations between dietary sodium and potassium intakes and mortality from cardiovascular disease: the Japan Collaborative Cohort Study for Evaluation of Cancer Risks. *Am J Clin Nutr* 2008 July;88(1):195-202.
- (38) Cook NR, Obarzanek E, Cutler JA et al. Joint effects of sodium and potassium intake on subsequent cardiovascular disease: the Trials of Hypertension Prevention follow-up study. *Arch Intern Med* 2009 January 12;169(1):32-40.
- (39) Yang Q, Liu T, Kuklina EV et al. Sodium and potassium intake and mortality among US adults: prospective data from the Third National Health and Nutrition Examination Survey. *Arch Intern Med* 2011 July 11;171(13):1183-91.
- (40) Chang HY, Hu YW, Yue CS et al. Effect of potassium-enriched salt on cardiovascular mortality and medical expenses of elderly men. *Am J Clin Nutr* 2006 June;83(6):1289-96.
- (41) Karppanen H, Mervaala E. Adherence to and population impact of non-pharmacological and pharmacological antihypertensive therapy. *J Hum Hypertens* 1996 February;10 Suppl 1:S57-S61.
- (42) Reimers CD, Knapp G, Reimers AK. Exercise as stroke prophylaxis. *Dtsch Arztebl Int* 2009 October;106(44):715-21.
- (43) Larson EB, Wang L, Bowen JD et al. Exercise is associated with reduced risk for incident dementia among persons 65 years of age and older. *Ann Intern Med* 2006 January 17;144(2):73-81.
- (44) Sumic A, Michael YL, Carlson NE, Howieson DB, Kaye JA. Physical activity and the risk of dementia in oldest old. *J Aging Health* 2007 April;19(2):242-59.
- (45) Andel R, Crowe M, Pedersen NL, Fratiglioni L, Johansson B, Gatz M. Physical exercise at midlife and risk of dementia three decades later: a population-based study of Swedish twins. *J Gerontol A Biol Sci Med Sci* 2008 January;63(1):62-6.

- (46) Jedrzejewski MK, Ewbank DC, Wang H, Trojanowski JQ. Exercise and cognition: results from the National Long Term Care Survey. *Alzheimers Dement* 2010 November;6(6):448-55.
- (47) Ahlskog JE, Geda YE, Graff-Radford NR, Petersen RC. Physical exercise as a preventive or disease-modifying treatment of dementia and brain aging. *Mayo Clin Proc* 2011 September;86(9):876-84.
- (48) Buchman AS, Boyle PA, Yu L, Shah RC, Wilson RS, Bennett DA. Total daily physical activity and the risk of AD and cognitive decline in older adults. *Neurology* 2012 April 24;78(17):1323-9.
- (49) Sofi F, Valecchi D, Bacci D et al. Physical activity and risk of cognitive decline: a meta-analysis of prospective studies. *J Intern Med* 2011 January;269(1):107-17.
- (50) Lange-Asschenfeldt C, Kojda G. Alzheimer's disease, cerebrovascular dysfunction and the benefits of exercise: from vessels to neurons. *Exp Gerontol* 2008 June;43(6):499-504.
- (51) Barrientos RM, Frank MG, Crysdale NY et al. Little exercise, big effects: reversing aging and infection-induced memory deficits, and underlying processes. *J Neurosci* 2011 August 10;31(32):11578-86.
- (52) Aguiar AS, Jr., Castro AA, Moreira EL et al. Short bouts of mild-intensity physical exercise improve spatial learning and memory in aging rats: involvement of hippocampal plasticity via AKT, CREB and BDNF signaling. *Mech Ageing Dev* 2011 November;132(11-12):560-7.
- (53) Rothman SM, Griffioen KJ, Wan R, Mattson MP. Brain-derived neurotrophic factor as a regulator of systemic and brain energy metabolism and cardiovascular health. *Ann N Y Acad Sci* 2012 August;1264(1):49-63.
- (54) Intlekofer KA, Cotman CW. Exercise counteracts declining hippocampal function in aging and Alzheimer's disease. *Neurobiol Dis* 2012 June 30.
- (55) Erickson KI, Voss MW, Prakash RS et al. Exercise training increases size of hippocampus and improves memory. *Proc Natl Acad Sci U S A* 2011 February 15;108(7):3017-22.
- (56) Adlard PA, Perreau VM, Pop V, Cotman CW. Voluntary exercise decreases amyloid load in a transgenic model of Alzheimer's disease. *J Neurosci* 2005 April 27;25(17):4217-21.
- (57) Parachikova A, Nichol KE, Cotman CW. Short-term exercise in aged Tg2576 mice alters neuroinflammation and improves cognition. *Neurobiol Dis* 2008 April;30(1):121-9.
- (58) Um HS, Kang EB, Leem YH et al. Exercise training acts as a therapeutic strategy for reduction of the pathogenic phenotypes for Alzheimer's disease in an NSE/APPsw-transgenic model. *Int J Mol Med* 2008 October;22(4):529-39.
- (59) Yuede CM, Zimmerman SD, Dong H et al. Effects of voluntary and forced exercise on plaque deposition, hippocampal volume, and behavior in the Tg2576 mouse model of Alzheimer's disease. *Neurobiol Dis* 2009 September;35(3):426-32.
- (60) Liu HL, Zhao G, Cai K, Zhao HH, Shi LD. Treadmill exercise prevents decline in spatial learning and memory in APP/PS1 transgenic mice through improvement of hippocampal long-term potentiation. *Behav Brain Res* 2011 April 15;218(2):308-14.

- (61) Garcia-Mesa Y, Lopez-Ramos JC, Gimenez-Llort L et al. Physical exercise protects against Alzheimer's disease in 3xTg-AD mice. *J Alzheimers Dis* 2011;24(3):421-54.
- (62) Kohman RA, Deyoung EK, Bhattacharya TK, Peterson LN, Rhodes JS. Wheel running attenuates microglia proliferation and increases expression of a proneurogenic phenotype in the hippocampus of aged mice. *Brain Behav Immun* 2012 July;26(5):803-10.
- (63) Soto ME, Secher M, Gillette-Guyonnet S et al. Weight loss and rapid cognitive decline in community-dwelling patients with Alzheimer's disease. *J Alzheimers Dis* 2012;28(3):647-54.
- (64) McCarty MF. Dietary saturate/unsaturate ratio as a determinant of adiposity. *Med Hypotheses* 2010 July;75(1):14-6.
- (65) Luchsinger JA, Gustafson DR. Adiposity and Alzheimer's disease. *Curr Opin Clin Nutr Metab Care* 2009 January;12(1):15-21.
- (66) Luchsinger JA. Diabetes, related conditions, and dementia. *J Neurol Sci* 2010 December 15;299(1-2):35-8.
- (67) Vanhanen M, Koivisto K, Moilanen L et al. Association of metabolic syndrome with Alzheimer disease: a population-based study. *Neurology* 2006 September 12;67(5):843-7.
- (68) Raffaitin C, Gin H, Empana JP et al. Metabolic syndrome and risk for incident Alzheimer's disease or vascular dementia: the Three-City Study. *Diabetes Care* 2009 January;32(1):169-74.
- (69) Profenno LA, Porsteinsson AP, Faraone SV. Meta-analysis of Alzheimer's disease risk with obesity, diabetes, and related disorders. *Biol Psychiatry* 2010 March 15;67(6):505-12.
- (70) Panza F, Frisardi V, Capurso C et al. Metabolic syndrome and cognitive impairment: current epidemiology and possible underlying mechanisms. *J Alzheimers Dis* 2010;21(3):691-724.
- (71) Forti P, Pisacane N, Rietti E et al. Metabolic syndrome and risk of dementia in older adults. *J Am Geriatr Soc* 2010 March;58(3):487-92.
- (72) Inelmen EM, Sergi G, Coin A, Girardi A, Manzato E. An open-ended question: Alzheimer's disease and involuntary weight loss: which comes first? *Aging Clin Exp Res* 2010 June;22(3):192-7.
- (73) Sims-Robinson C, Kim B, Rosko A, Feldman EL. How does diabetes accelerate Alzheimer disease pathology? *Nat Rev Neurol* 2010 October;6(10):551-9.
- (74) Exalto LG, Whitmer RA, Kappelle LJ, Biessels GJ. An update on type 2 diabetes, vascular dementia and Alzheimer's disease. *Exp Gerontol* 2012 August 3.
- (75) Brothers HM, Marchalant Y, Wenk GL. Caffeine attenuates lipopolysaccharide-induced neuroinflammation. *Neurosci Lett* 2010 August 16;480(2):97-100.
- (76) Takahashi RN, Pamplona FA, Prediger RD. Adenosine receptor antagonists for cognitive dysfunction: a review of animal studies. *Front Biosci* 2008;13:2614-32.

- (77) Dall'Igna OP, Porciuncula LO, Souza DO, Cunha RA, Lara DR. Neuroprotection by caffeine and adenosine A2A receptor blockade of beta-amyloid neurotoxicity. *Br J Pharmacol* 2003 April;138(7):1207-9.
- (78) Arendash GW, Schleif W, Rezai-Zadeh K et al. Caffeine protects Alzheimer's mice against cognitive impairment and reduces brain beta-amyloid production. *Neuroscience* 2006 November 3;142(4):941-52.
- (79) Cao C, Cirrito JR, Lin X et al. Caffeine suppresses amyloid-beta levels in plasma and brain of Alzheimer's disease transgenic mice. *J Alzheimers Dis* 2009;17(3):681-97.
- (80) Dall'Igna OP, Porciuncula LO, Souza DO, Cunha RA, Lara DR. Neuroprotection by caffeine and adenosine A2A receptor blockade of beta-amyloid neurotoxicity. *Br J Pharmacol* 2003 April;138(7):1207-9.
- (81) Canas PM, Porciuncula LO, Cunha GM et al. Adenosine A2A receptor blockade prevents synaptotoxicity and memory dysfunction caused by beta-amyloid peptides via p38 mitogen-activated protein kinase pathway. *J Neurosci* 2009 November 25;29(47):14741-51.
- (82) Cunha RA, Agostinho PM. Chronic caffeine consumption prevents memory disturbance in different animal models of memory decline. *J Alzheimers Dis* 2010;20 Suppl 1:S95-116.
- (83) Maia L, de MA. Does caffeine intake protect from Alzheimer's disease? *Eur J Neurol* 2002 July;9(4):377-82.
- (84) Barranco Quintana JL, Allam MF, Serrano Del CA, Fernandez-Crehuet NR. Alzheimer's disease and coffee: a quantitative review. *Neurol Res* 2007 January;29(1):91-5.
- (85) Ritchie K, Carriere I, de MA et al. The neuroprotective effects of caffeine: a prospective population study (the Three City Study). *Neurology* 2007 August 7;69(6):536-45.
- (86) Eskelinen MH, Ngandu T, Tuomilehto J, Soininen H, Kivipelto M. Midlife coffee and tea drinking and the risk of late-life dementia: a population-based CAIDE study. *J Alzheimers Dis* 2009;16(1):85-91.
- (87) Santos C, Costa J, Santos J, Vaz-Carneiro A, Lunet N. Caffeine intake and dementia: systematic review and meta-analysis. *J Alzheimers Dis* 2010;20 Suppl 1:S187-S204.
- (88) Santos C, Lunet N, Azevedo A, de MA, Ritchie K, Barros H. Caffeine intake is associated with a lower risk of cognitive decline: a cohort study from Portugal. *J Alzheimers Dis* 2010;20 Suppl 1:S175-S185.
- (89) Eskelinen MH, Kivipelto M. Caffeine as a protective factor in dementia and Alzheimer's disease. *J Alzheimers Dis* 2010;20 Suppl 1:S167-S174.
- (90) Laitala VS, Kaprio J, Koskenvuo M, Raiha I, Rinne JO, Silventoinen K. Coffee drinking in middle age is not associated with cognitive performance in old age. *Am J Clin Nutr* 2009 September;90(3):640-6.
- (91) Gelber RP, Petrovitch H, Masaki KH, Ross GW, White LR. Coffee intake in midlife and risk of dementia and its neuropathologic correlates. *J Alzheimers Dis* 2011;23(4):607-15.

- (92) Cao C, Loewenstein DA, Lin X et al. High Blood caffeine levels in MCI linked to lack of progression to dementia. *J Alzheimers Dis* 2012;30(3):559-72.
- (93) Freedman ND, Park Y, Abnet CC, Hollenbeck AR, Sinha R. Association of coffee drinking with total and cause-specific mortality. *N Engl J Med* 2012 May 17;366(20):1891-904.
- (94) Matos M, Augusto E, Santos-Rodrigues AD et al. Adenosine A(2A) receptors modulate glutamate uptake in cultured astrocytes and gliosomes. *Glia* 2012 May;60(5):702-16.
- (95) Matos M, Augusto E, Machado NJ, Dos Santos-Rodrigues A, Cunha RA, Agostinho P. Astrocytic Adenosine A2A Receptors Control the Amyloid-beta Peptide-Induced Decrease of Glutamate Uptake. *J Alzheimers Dis* 2012 January 1;31(3):555-67.
- (96) Xu J, Kurup P, Zhang Y et al. Extrasynaptic NMDA receptors couple preferentially to excitotoxicity via calpain-mediated cleavage of STEP. *J Neurosci* 2009 July 22;29(29):9330-43.
- (97) Hardingham GE, Bading H. Synaptic versus extrasynaptic NMDA receptor signalling: implications for neurodegenerative disorders. *Nat Rev Neurosci* 2010 October;11(10):682-96.
- (98) Li S, Jin M, Koeglsperger T, Shepardson NE, Shankar GM, Selkoe DJ. Soluble Aβ oligomers inhibit long-term potentiation through a mechanism involving excessive activation of extrasynaptic NR2B-containing NMDA receptors. *J Neurosci* 2011 May 4;31(18):6627-38.
- (99) Wurtman RJ, Ulus IH, Cansev M, Watkins CJ, Wang L, Marzloff G. Synaptic proteins and phospholipids are increased in gerbil brain by administering uridine plus docosahexaenoic acid orally. *Brain Res* 2006 May 9;1088(1):83-92.
- (100) Luchtman DW, Song C. Cognitive enhancement by omega-3 fatty acids from child-hood to old age: Findings from animal and clinical studies. *Neuropharmacology* 2012 July 27.
- (101) Labrousse VF, Nadjar A, Joffre C et al. Short-term long chain omega3 diet protects from neuroinflammatory processes and memory impairment in aged mice. *PLoS ONE* 2012;7(5):e36861.
- (102) Ueda Y, Wang MF, Irei AV, Sarukura N, Sakai T, Hsu TF. Effect of dietary lipids on longevity and memory in the SAMP8 mice. *J Nutr Sci Vitaminol (Tokyo)* 2011;57(1):36-41.
- (103) Hooijmans CR, Van der Zee CE, Dederen PJ et al. DHA and cholesterol containing diets influence Alzheimer-like pathology, cognition and cerebral vasculature in APP^{swe}/PS1^{dE9} mice. *Neurobiol Dis* 2009 March;33(3):482-98.
- (104) Petursdottir AL, Farr SA, Morley JE, Banks WA, Skuladottir GV. Effect of dietary n-3 polyunsaturated fatty acids on brain lipid fatty acid composition, learning ability, and memory of senescence-accelerated mouse. *J Gerontol A Biol Sci Med Sci* 2008 November;63(11):1153-60.
- (105) Gamoh S, Hashimoto M, Hossain S, Masumura S. Chronic administration of docosahexaenoic acid improves the performance of radial arm maze task in aged rats. *Clin Exp Pharmacol Physiol* 2001 April;28(4):266-70.

- (106) Arendash GW, Jensen MT, Salem N, Jr. et al. A diet high in omega-3 fatty acids does not improve or protect cognitive performance in Alzheimer's transgenic mice. *Neuroscience* 2007 October 26;149(2):286-302.
- (107) Oster T, Pillot T. Docosahexaenoic acid and synaptic protection in Alzheimer's disease mice. *Biochim Biophys Acta* 2010 August;1801(8):791-8.
- (108) van Gelder BM, Tijhuis M, Kalmijn S, Kromhout D. Fish consumption, n-3 fatty acids, and subsequent 5-y cognitive decline in elderly men: the Zutphen Elderly Study. *Am J Clin Nutr* 2007 April;85(4):1142-7.
- (109) Yurko-Mauro K. Cognitive and cardiovascular benefits of docosahexaenoic acid in aging and cognitive decline. *Curr Alzheimer Res* 2010 May;7(3):190-6.
- (110) van Gelder BM, Tijhuis M, Kalmijn S, Kromhout D. Fish consumption, n-3 fatty acids, and subsequent 5-y cognitive decline in elderly men: the Zutphen Elderly Study. *Am J Clin Nutr* 2007 April;85(4):1142-7.
- (111) Kalmijn S, Feskens EJ, Launer LJ, Kromhout D. Polyunsaturated fatty acids, antioxidants, and cognitive function in very old men. *Am J Epidemiol* 1997 January 1;145(1):33-41.
- (112) Morris MC, Evans DA, Tangney CC, Bienias JL, Wilson RS. Fish consumption and cognitive decline with age in a large community study. *Arch Neurol* 2005 December;62(12):1849-53.
- (113) Morris MC, Evans DA, Bienias JL et al. Consumption of fish and n-3 fatty acids and risk of incident Alzheimer disease. *Arch Neurol* 2003 July;60(7):940-6.
- (114) Yurko-Mauro K, McCarthy D, Rom D et al. Beneficial effects of docosahexaenoic acid on cognition in age-related cognitive decline. *Alzheimers Dement* 2010 November;6(6):456-64.
- (115) Quinn JF, Raman R, Thomas RG et al. Docosahexaenoic acid supplementation and cognitive decline in Alzheimer disease: a randomized trial. *JAMA* 2010 November 3;304(17):1903-11.
- (116) Lemke SL, Vicini JL, Su H et al. Dietary intake of stearidonic acid-enriched soybean oil increases the omega-3 index: randomized, double-blind clinical study of efficacy and safety. *Am J Clin Nutr* 2010 October;92(4):766-75.
- (117) Neveu I, Naveilhan P, Mena C, Wion D, Brachet P, Garabedian M. Synthesis of 1,25-dihydroxyvitamin D3 by rat brain macrophages in vitro. *J Neurosci Res* 1994 June 1;38(2):214-20.
- (118) Lefebvre dC, Montero-Menei CN, Bernard R, Couez D. Vitamin D3 inhibits proinflammatory cytokines and nitric oxide production by the EOC13 microglial cell line. *J Neurosci Res* 2003 February 15;71(4):575-82.
- (119) Garcion E, Nataf S, Berod A, Darcy F, Brachet P. 1,25-Dihydroxyvitamin D3 inhibits the expression of inducible nitric oxide synthase in rat central nervous system during experimental allergic encephalomyelitis. *Brain Res Mol Brain Res* 1997 May;45(2):255-67.
- (120) Moore ME, Piazza A, McCartney Y, Lynch MA. Evidence that vitamin D3 reverses age-related inflammatory changes in the rat hippocampus. *Biochem Soc Trans* 2005 August;33(Pt 4):573-7.

- (121) Oudshoorn C, Mattace-Raso FU, Van d, V, Colin EM, van der Cammen TJ. Higher serum vitamin D3 levels are associated with better cognitive test performance in patients with Alzheimer's disease. *Dement Geriatr Cogn Disord* 2008;25(6):539-43.
- (122) Llewellyn DJ, Lang IA, Langa KM et al. Vitamin D and risk of cognitive decline in elderly persons. *Arch Intern Med* 2010 July 12;170(13):1135-41.
- (123) Breitling LP, Perna L, Muller H, Raum E, Kliegel M, Brenner H. Vitamin D and cognitive functioning in the elderly population in Germany. *Exp Gerontol* 2012 January;47(1):122-7.
- (124) Annweiler C, Beauchet O. Vitamin D-mentia: randomized clinical trials should be the next step. *Neuroepidemiology* 2011;37(3-4):249-58.
- (125) Skalska A, Galas A, Grodzicki T. 25-hydroxyvitamin D and physical and cognitive performance in older people with chronic conditions. *Pol Arch Med Wewn* 2012;122(4):162-9.
- (126) Slinin Y, Paudel M, Taylor BC et al. Association Between Serum 25(OH) Vitamin D and the Risk of Cognitive Decline in Older Women. *J Gerontol A Biol Sci Med Sci* 2012 March 27.
- (127) Annweiler C, Rolland Y, Schott AM et al. Higher Vitamin D Dietary Intake Is Associated With Lower Risk of Alzheimer's Disease: A 7-Year Follow-up. *J Gerontol A Biol Sci Med Sci* 2012 April 13.
- (128) Soni M, Kos K, Lang IA, Jones K, Melzer D, Llewellyn DJ. Vitamin D and cognitive function. *Scand J Clin Lab Invest Suppl* 2012 April;243:79-82.
- (129) Holick MF. Vitamin D: a D-Lightful health perspective. *Nutr Rev* 2008 October;66(10 Suppl 2):S182-S194.
- (130) Vieth R. Vitamin D supplementation, 25-hydroxyvitamin D concentrations, and safety. *Am J Clin Nutr* 1999 May;69(5):842-56.
- (131) Pawate S, Shen Q, Fan F, Bhat NR. Redox regulation of glial inflammatory response to lipopolysaccharide and interferongamma. *J Neurosci Res* 2004 August 15;77(4):540-51.
- (132) Choi SH, Aid S, Kim HW, Jackson SH, Bosetti F. Inhibition of NADPH oxidase promotes alternative and anti-inflammatory microglial activation during neuroinflammation. *J Neurochem* 2012 January;120(2):292-301.
- (133) Koppula S, Kumar H, Kim IS, Choi DK. Reactive oxygen species and inhibitors of inflammatory enzymes, NADPH oxidase, and iNOS in experimental models of Parkinson's disease. *Mediators Inflamm* 2012;2012:823902.
- (134) Peterson LJ, Flood PM. Oxidative stress and microglial cells in Parkinson's disease. *Mediators Inflamm* 2012;2012:401264.
- (135) Block ML. NADPH oxidase as a therapeutic target in Alzheimer's disease. *BMC Neurosci* 2008;9 Suppl 2:S8.
- (136) Surace MJ, Block ML. Targeting microglia-mediated neurotoxicity: the potential of NOX2 inhibitors. *Cell Mol Life Sci* 2012 July;69(14):2409-27.

- (137) Ansari MA, Scheff SW. NADPH-oxidase activation and cognition in Alzheimer disease progression. *Free Radic Biol Med* 2011 July 1;51(1):171-8.
- (138) Mrak RE, Griffin WS. Dementia with Lewy bodies: Definition, diagnosis, and pathogenic relationship to Alzheimer's disease. *Neuropsychiatr Dis Treat* 2007;3(5):619-25.
- (139) Serrano F, Chang A, Hernandez C, Pautler RG, Sweatt JD, Klann E. NADPH oxidase mediates beta-amyloid peptide-induced activation of ERK in hippocampal organotypic cultures. *Mol Brain* 2009;2:31.
- (140) He Y, Cui J, Lee JC et al. Prolonged exposure of cortical neurons to oligomeric amyloid-beta impairs NMDA receptor function via NADPH oxidase-mediated ROS production: protective effect of green tea (-)-epigallocatechin-3-gallate. *ASN Neuro* 2011;3(1):e00050.
- (141) Du H, Guo L, Fang F et al. Cyclophilin D deficiency attenuates mitochondrial and neuronal perturbation and ameliorates learning and memory in Alzheimer's disease. *Nat Med* 2008 October;14(10):1097-105.
- (142) Massaad CA, Washington TM, Pautler RG, Klann E. Overexpression of SOD-2 reduces hippocampal superoxide and prevents memory deficits in a mouse model of Alzheimer's disease. *Proc Natl Acad Sci U S A* 2009 August 11;106(32):13576-81.
- (143) Massaad CA, Pautler RG, Klann E. Mitochondrial superoxide: a key player in Alzheimer's disease. *Aging (Albany NY)* 2009 September;1(9):758-61.
- (144) McManus MJ, Murphy MP, Franklin JL. The mitochondria-targeted antioxidant MitoQ prevents loss of spatial memory retention and early neuropathology in a transgenic mouse model of Alzheimer's disease. *J Neurosci* 2011 November 2;31(44):15703-15.
- (145) Dugan LL, Ali SS, Shekhtman G et al. IL-6 mediated degeneration of forebrain GABAergic interneurons and cognitive impairment in aged mice through activation of neuronal NADPH oxidase. *PLoS ONE* 2009;4(5):e5518.
- (146) Stanley EM, Fadel JR, Mott DD. Interneuron loss reduces dendritic inhibition and GABA release in hippocampus of aged rats. *Neurobiol Aging* 2012 February;33(2):431-13.
- (147) Park L, Anrather J, Zhou P et al. NADPH-oxidase-derived reactive oxygen species mediate the cerebrovascular dysfunction induced by the amyloid beta peptide. *J Neurosci* 2005 February 16;25(7):1769-77.
- (148) Park L, Zhou P, Pitstick R et al. Nox2-derived radicals contribute to neurovascular and behavioral dysfunction in mice overexpressing the amyloid precursor protein. *Proc Natl Acad Sci U S A* 2008 January 29;105(4):1347-52.
- (149) Ito D, Murata M, Watanabe K et al. C242T polymorphism of NADPH oxidase p22 PHOX gene and ischemic cerebrovascular disease in the Japanese population. *Stroke* 2000 April;31(4):936-9.
- (150) Genius J, Grau AJ, Lichy C. The C242T polymorphism of the NAD(P)H oxidase p22phox subunit is associated with an enhanced risk for cerebrovascular disease at a young age. *Cerebrovasc Dis* 2008;26(4):430-3.

- (151) Tang XN, Cairns B, Kim JY, Yenari MA. NADPH oxidase in stroke and cerebrovascular disease. *Neurol Res* 2012 May;34(4):338-45.
- (152) Kahles T, Brandes RP. Which NADPH Oxidase Isoform Is Relevant for Ischemic Stroke? The Case for Nox 2. *Antioxid Redox Signal* 2012 August 20.
- (153) Kim HA, Miller AA, Drummond GR et al. Vascular cognitive impairment and Alzheimer's disease: role of cerebral hypoperfusion and oxidative stress. *Naunyn Schmiedebergs Arch Pharmacol* 2012 August 8.
- (154) Lanone S, Bloc S, Foresti R et al. Bilirubin decreases nos2 expression via inhibition of NAD(P)H oxidase: implications for protection against endotoxic shock in rats. *FASEB J* 2005 November;19(13):1890-2.
- (155) Matsumoto H, Ishikawa K, Itabe H, Maruyama Y. Carbon monoxide and bilirubin from heme oxygenase-1 suppresses reactive oxygen species generation and plasminogen activator inhibitor-1 induction. *Mol Cell Biochem* 2006 October;291(1-2):21-8.
- (156) Jiang F, Roberts SJ, Datla S, Dusting GJ. NO modulates NADPH oxidase function via heme oxygenase-1 in human endothelial cells. *Hypertension* 2006 November;48(5):950-7.
- (157) Datla SR, Dusting GJ, Mori TA, Taylor CJ, Croft KD, Jiang F. Induction of heme oxygenase-1 in vivo suppresses NADPH oxidase derived oxidative stress. *Hypertension* 2007 October;50(4):636-42.
- (158) Terry MJ, Maines MD, Lagarias JC. Inactivation of phytochrome- and phycobiliprotein-chromophore precursors by rat liver biliverdin reductase. *J Biol Chem* 1993 December 15;268(35):26099-106.
- (159) McCarty MF. Clinical potential of Spirulina as a source of phycocyanobilin. *J Med Food* 2007 December;10(4):566-70.
- (160) Zheng J, Inoguchi T, Sasaki S et al. Phycocyanin and phycocyanobilin from *Spirulina platensis* protect against diabetic nephropathy by inhibiting oxidative stress. *Am J Physiol Regul Integr Comp Physiol* 2013 January 15;304(2):R110-R120.
- (161) Romay C, Gonzalez R, Ledon N, Ramirez D, Rimbau V. C-phycocyanin: a biliprotein with antioxidant, anti-inflammatory and neuroprotective effects. *Curr Protein Pept Sci* 2003 June;4(3):207-16.
- (162) Chamorro G, Perez-Albiter M, Serrano-Garcia N, Mares-Samano JJ, Rojas P. *Spirulina maxima* pretreatment partially protects against 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine neurotoxicity. *Nutr Neurosci* 2006 October;9(5-6):207-12.
- (163) Rimbau V, Camins A, Romay C, Gonzalez R, Pallas M. Protective effects of C-phycocyanin against kainic acid-induced neuronal damage in rat hippocampus. *Neurosci Lett* 1999 December 3;276(2):75-8.
- (164) Pabon MM, Jernberg JN, Morganti J et al. A *Spirulina*-Enhanced Diet Provides Neuroprotection in an alpha-Synuclein Model of Parkinson's Disease. *PLoS ONE* 2012;7(9):e45256.

- (165) Hwang JH, Lee IT, Jeng KC et al. Spirulina prevents memory dysfunction, reduces oxidative stress damage and augments antioxidant activity in senescence-accelerated mice. *J Nutr Sci Vitaminol (Tokyo)* 2011;57(2):186-91.
- (166) McCarty MF, Barroso-Aranda J, Contreras F. Oral phycocyanobilin may diminish the pathogenicity of activated brain microglia in neurodegenerative disorders. *Med Hypotheses* 2010 March;74(3):601-5.
- (167) Penton-Rol G, Marin-Prida J, Pardo-Andreu G et al. C-Phycocyanin is neuroprotective against global cerebral ischemia/reperfusion injury in gerbils. *Brain Res Bull* 2011 August 10;86(1-2):42-52.
- (168) Palozza P, Krinsky NI. Astaxanthin and canthaxanthin are potent antioxidants in a membrane model. *Arch Biochem Biophys* 1992 September;297(2):291-5.
- (169) Naguib YM. Antioxidant activities of astaxanthin and related carotenoids. *J Agric Food Chem* 2000 April;48(4):1150-4.
- (170) Liu X, Shibata T, Hisaka S, Osawa T. Astaxanthin inhibits reactive oxygen species-mediated cellular toxicity in dopaminergic SH-SY5Y cells via mitochondria-targeted protective mechanism. *Brain Res* 2009 February 13;1254:18-27.
- (171) Wolf AM, Asoh S, Hiranuma H et al. Astaxanthin protects mitochondrial redox state and functional integrity against oxidative stress. *J Nutr Biochem* 2010 May;21(5):381-9.
- (172) Lee DH, Kim CS, Lee YJ. Astaxanthin protects against MPTP/MPP+-induced mitochondrial dysfunction and ROS production in vivo and in vitro. *Food Chem Toxicol* 2010 November 4.
- (173) Shen H, Kuo CC, Chou J et al. Astaxanthin reduces ischemic brain injury in adult rats. *FASEB J* 2009 June;23(6):1958-68.
- (174) Pashkow FJ, Watumull DG, Campbell CL. Astaxanthin: a novel potential treatment for oxidative stress and inflammation in cardiovascular disease. *Am J Cardiol* 2008 May 22;101(10A):58D-68D.
- (175) Liu X, Osawa T. Astaxanthin protects neuronal cells against oxidative damage and is a potent candidate for brain food. *Forum Nutr* 2009;61:129-35.
- (176) Kim YH, Koh HK, Kim DS. Down-regulation of IL-6 production by astaxanthin via ERK-, MSK-, and NF-kappaB-mediated signals in activated microglia. *Int Immunopharmacol* 2010 December;10(12):1560-72.
- (177) Choi SK, Park YS, Choi DK, Chang HI. Effects of astaxanthin on the production of NO and the expression of COX-2 and iNOS in LPS-stimulated BV2 microglial cells. *J Microbiol Biotechnol* 2008 December;18(12):1990-6.
- (178) Karppi J, Rissanen TH, Nyyssonen K et al. Effects of astaxanthin supplementation on lipid peroxidation. *Int J Vitam Nutr Res* 2007 January;77(1):3-11.
- (179) Park JS, Chyun JH, Kim YK, Line LL, Chew BP. Astaxanthin decreased oxidative stress and inflammation and enhanced immune response in humans. *Nutr Metab (Lond)* 2010;7:18.

- (180) Kim JH, Chang MJ, Choi HD et al. Protective effects of Haematococcus astaxanthin on oxidative stress in healthy smokers. *J Med Food* 2011 November;14(11):1469-75.
- (181) Yoshida H, Yanai H, Ito K et al. Administration of natural astaxanthin increases serum HDL-cholesterol and adiponectin in subjects with mild hyperlipidemia. *Atherosclerosis* 2010 April;209(2):520-3.
- (182) Comhaire FH, El GY, Mahmoud A, Eertmans F, Schoonjans F. Combined conventional/antioxidant "Astaxanthin" treatment for male infertility: a double blind, randomized trial. *Asian J Androl* 2005 September;7(3):257-62.
- (183) Sano M, Ernesto C, Thomas RG et al. A controlled trial of selegiline, alpha-tocopherol, or both as treatment for Alzheimer's disease. The Alzheimer's Disease Cooperative Study. *N Engl J Med* 1997 April 24;336(17):1216-22.
- (184) Talalay P, Fahey JW. Phytochemicals from cruciferous plants protect against cancer by modulating carcinogen metabolism. *J Nutr* 2001 November;131(11 Suppl):3027S-33S.
- (185) Na HK, Surh YJ. Modulation of Nrf2-mediated antioxidant and detoxifying enzyme induction by the green tea polyphenol EGCG. *Food Chem Toxicol* 2008 April;46(4):1271-8.
- (186) Chen C, Pung D, Leong V et al. Induction of detoxifying enzymes by garlic organosulfur compounds through transcription factor Nrf2: effect of chemical structure and stress signals. *Free Radic Biol Med* 2004 November 15;37(10):1578-90.
- (187) Flier J, Van Muiswinkel FL, Jongenelen CA, Drukarch B. The neuroprotective antioxidant alpha-lipoic acid induces detoxication enzymes in cultured astroglial cells. *Free Radic Res* 2002 June;36(6):695-9.
- (188) Cao Z, Tsang M, Zhao H, Li Y. Induction of endogenous antioxidants and phase 2 enzymes by alpha-lipoic acid in rat cardiac H9C2 cells: protection against oxidative injury. *Biochem Biophys Res Commun* 2003 October 24;310(3):979-85.
- (189) Jia Z, Hallur S, Zhu H, Li Y, Misra HP. Potent upregulation of glutathione and NAD(P)H:quinone oxidoreductase 1 by alpha-lipoic acid in human neuroblastoma SH-SY5Y cells: protection against neurotoxicant-elicited cytotoxicity. *Neurochem Res* 2008 May;33(5):790-800.
- (190) Packer L, Tritschler HJ, Wessel K. Neuroprotection by the metabolic antioxidant alpha-lipoic acid. *Free Radic Biol Med* 1997;22(1-2):359-78.
- (191) Ziegler D, Ametov A, Barinov A et al. Oral treatment with alpha-lipoic acid improves symptomatic diabetic polyneuropathy: the SYDNEY 2 trial. *Diabetes Care* 2006 November;29(11):2365-70.
- (192) Surh YJ, Kundu JK, Na HK. Nrf2 as a master redox switch in turning on the cellular signaling involved in the induction of cytoprotective genes by some chemopreventive phytochemicals. *Planta Med* 2008 October;74(13):1526-39.

- (193) Zhang DD, Lo SC, Cross JV, Templeton DJ, Hannink M. Keap1 is a redox-regulated substrate adaptor protein for a Cul3-dependent ubiquitin ligase complex. *Mol Cell Biol* 2004 December;24(24):10941-53.
- (194) Li W, Thakor N, Xu EY et al. An internal ribosomal entry site mediates redox-sensitive translation of Nrf2. *Nucleic Acids Res* 2010 January;38(3):778-88.
- (195) Shay KP, Michels AJ, Li W, Kong AN, Hagen TM. Cap-independent Nrf2 translation is part of a lipoic acid-stimulated detoxification stress response. *Biochim Biophys Acta* 2012 June;1823(6):1102-9.
- (196) Kanninen K, Heikkinen R, Malm T et al. Intrahippocampal injection of a lentiviral vector expressing Nrf2 improves spatial learning in a mouse model of Alzheimer's disease. *Proc Natl Acad Sci U S A* 2009 September 22;106(38):16505-10.
- (197) Yang YC, Lu FH, Wu JS, Wu CH, Chang CJ. The protective effect of habitual tea consumption on hypertension. *Arch Intern Med* 2004 July 26;164(14):1534-40.
- (198) Jesudason EP, Masilamoni JG, Jesudoss KS, Jayakumar R. The protective role of DL-alpha-lipoic acid in the oxidative vulnerability triggered by Abeta-amyloid vaccination in mice. *Mol Cell Biochem* 2005 February;270(1-2):29-37.
- (199) Quinn JF, Bussiere JR, Hammond RS et al. Chronic dietary alpha-lipoic acid reduces deficits in hippocampal memory of aged Tg2576 mice. *Neurobiol Aging* 2007 February;28(2):213-25.
- (200) Hager K, Kenklies M, McAfoose J, Engel J, Munch G. Alpha-lipoic acid as a new treatment option for Alzheimer's disease--a 48 months follow-up analysis. *J Neural Transm Suppl* 2007;(72):189-93.
- (201) Maczurek A, Hager K, Kenklies M et al. Lipoic acid as an anti-inflammatory and neuroprotective treatment for Alzheimer's disease. *Adv Drug Deliv Rev* 2008 October;60(13-14):1463-70.
- (202) Rodriguez C, Mayo JC, Sainz RM et al. Regulation of antioxidant enzymes: a significant role for melatonin. *J Pineal Res* 2004 January;36(1):1-9.
- (203) Wang JZ, Wang ZF. Role of melatonin in Alzheimer-like neurodegeneration. *Acta Pharmacol Sin* 2006 January;27(1):41-9.
- (204) Cheng Y, Feng Z, Zhang QZ, Zhang JT. Beneficial effects of melatonin in experimental models of Alzheimer disease. *Acta Pharmacol Sin* 2006 February;27(2):129-39.
- (205) Shen Y, Zhang G, Liu L, Xu S. Suppressive effects of melatonin on amyloid-beta-induced glial activation in rat hippocampus. *Arch Med Res* 2007 April;38(3):284-90.
- (206) Zhou J, Zhang S, Zhao X, Wei T. Melatonin impairs NADPH oxidase assembly and decreases superoxide anion production in microglia exposed to amyloid-beta1-42. *J Pineal Res* 2008 September;45(2):157-65.
- (207) Ionov M, Burchell V, Klajnert B, Bryszewska M, Abramov AY. Mechanism of neuroprotection of melatonin against beta-amyloid neurotoxicity. *Neuroscience* 2011 April 28;180:229-37.

- (208) Dragicevic N, Copes N, O'Neal-Moffitt G et al. Melatonin treatment restores mitochondrial function in Alzheimer's mice: a mitochondrial protective role of melatonin membrane receptor signaling. *J Pineal Res* 2011 August;51(1):75-86.
- (209) Rosales-Corral SA, Acuna-Castroviejo D, Coto-Montes A et al. Alzheimer's disease: pathological mechanisms and the beneficial role of melatonin. *J Pineal Res* 2012 March;52(2):167-202.
- (210) Garcia-Mesa Y, Gimenez-Llort L, Lopez LC et al. Melatonin plus physical exercise are highly neuroprotective in the 3xTg-AD mouse. *Neurobiol Aging* 2012 June;33(6):1124-9.
- (211) Cardinali DP, Furio AM, Brusco LI. Clinical aspects of melatonin intervention in Alzheimer's disease progression. *Curr Neuropharmacol* 2010 September;8(3):218-27.
- (212) Atkuri KR, Mantovani JJ, Herzenberg LA, Herzenberg LA. N-Acetylcysteine--a safe antidote for cysteine/glutathione deficiency. *Curr Opin Pharmacol* 2007 August;7(4):355-9.
- (213) Dodd S, Dean O, Copolov DL, Malhi GS, Berk M. N-acetylcysteine for antioxidant therapy: pharmacology and clinical utility. *Expert Opin Biol Ther* 2008 December;8(12):1955-62.
- (214) Takagi Y, Kurihara S, Higashi N et al. Combined administration of (L)-cystine and (L)-theanine enhances immune functions and protects against influenza virus infection in aged mice. *J Vet Med Sci* 2010 February;72(2):157-65.
- (215) Kojima S, Miyazaki Y, Honda T, Kiyozumi M, Shimada H, Funakoshi T. Effect of L-cystine on toxicity of paraquat in mice. *Toxicol Lett* 1992 January;60(1):75-82.
- (216) Munoz AM, Rey P, Soto-Otero R, Guerra MJ, Labandeira-Garcia JL. Systemic administration of N-acetylcysteine protects dopaminergic neurons against 6-hydroxydopamine-induced degeneration. *J Neurosci Res* 2004 May 15;76(4):551-62.
- (217) Park SW, Kim SH, Park KH et al. Preventive effect of antioxidants in MPTP-induced mouse model of Parkinson's disease. *Neurosci Lett* 2004 June 17;363(3):243-6.
- (218) Arakawa M, Ito Y. N-acetylcysteine and neurodegenerative diseases: Basic and clinical pharmacology. *Cerebellum* 2007;6(4):308-14.
- (219) Clark J, Clore EL, Zheng K, Adame A, Masliah E, Simon DK. Oral N-acetyl-cysteine attenuates loss of dopaminergic terminals in alpha-synuclein overexpressing mice. *PLoS ONE* 2010;5(8):e12333.
- (220) Berman AE, Chan WY, Brennan AM et al. N-acetylcysteine prevents loss of dopaminergic neurons in the EAAC1^{-/-} mouse. *Ann Neurol* 2011 March;69(3):509-20.
- (221) Martinez-Banaclocha MA. N-acetyl-cysteine in the treatment of Parkinson's disease. What are we waiting for? *Med Hypotheses* 2012 July;79(1):8-12.
- (222) Fu AL, Dong ZH, Sun MJ. Protective effect of N-acetyl-L-cysteine on amyloid beta-peptide-induced learning and memory deficits in mice. *Brain Res* 2006 September 13;1109(1):201-6.

- (223) Hsiao YH, Kuo JR, Chen SH, Gean PW. Amelioration of social isolation-triggered onset of early Alzheimer's disease-related cognitive deficit by N-acetylcysteine in a transgenic mouse model. *Neurobiol Dis* 2012 March;45(3):1111-20.
- (224) Liberatore GT, Jackson-Lewis V, Vukosavic S et al. Inducible nitric oxide synthase stimulates dopaminergic neurodegeneration in the MPTP model of Parkinson disease. *Nat Med* 1999 December;5(12):1403-9.
- (225) Dehmer T, Lindenau J, Haid S, Dichgans J, Schulz JB. Deficiency of inducible nitric oxide synthase protects against MPTP toxicity in vivo. *J Neurochem* 2000 May;74(5):2213-6.
- (226) Wu DC, Jackson-Lewis V, Vila M et al. Blockade of microglial activation is neuroprotective in the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine mouse model of Parkinson disease. *J Neurosci* 2002 March 1;22(5):1763-71.
- (227) Tieu K, Ischiropoulos H, Przedborski S. Nitric oxide and reactive oxygen species in Parkinson's disease. *IUBMB Life* 2003 June;55(6):329-35.
- (228) Irvani MM, Haddon CO, Rose S, Jenner P. 3-Nitrotyrosine-dependent dopaminergic neurotoxicity following direct nigral administration of a peroxynitrite but not a nitric oxide donor. *Brain Res* 2006 January 5;1067(1):256-62.
- (229) Giasson BI, Duda JE, Murray IV et al. Oxidative damage linked to neurodegeneration by selective alpha-synuclein nitration in synucleinopathy lesions. *Science* 2000 November 3;290(5493):985-9.
- (230) Duda JE, Giasson BI, Chen Q et al. Widespread nitration of pathological inclusions in neurodegenerative synucleinopathies. *Am J Pathol* 2000 November;157(5):1439-45.
- (231) Danielson SR, Held JM, Schilling B, Oo M, Gibson BW, Andersen JK. Preferentially increased nitration of alpha-synuclein at tyrosine-39 in a cellular oxidative model of Parkinson's disease. *Anal Chem* 2009 September 15;81(18):7823-8.
- (232) Gao HM, Kotzbauer PT, Uryu K, Leight S, Trojanowski JQ, Lee VM. Neuroinflammation and oxidation/nitration of alpha-synuclein linked to dopaminergic neurodegeneration. *J Neurosci* 2008 July 23;28(30):7687-98.
- (233) Kirsch M, Lehnig M, Korth HG, Sustmann R, de GH. Inhibition of peroxynitrite-induced nitration of tyrosine by glutathione in the presence of carbon dioxide through both radical repair and peroxynitrate formation. *Chemistry* 2001 August 3;7(15):3313-20.
- (234) Jazwa A, Rojo AI, Innamorato NG, Hesse M, Fernandez-Ruiz J, Cuadrado A. Pharmacological targeting of the transcription factor Nrf2 at the basal ganglia provides disease modifying therapy for experimental parkinsonism. *Antioxid Redox Signal* 2011 June 15;14(12):2347-60.
- (235) Zhang H, Jia H, Liu J et al. Combined R-alpha-lipoic acid and acetyl-L-carnitine exerts efficient preventative effects in a cellular model of Parkinson's disease. *J Cell Mol Med* 2010 January;14(1-2):215-25.
- (236) Karunakaran S, Diwakar L, Saeed U et al. Activation of apoptosis signal regulating kinase 1 (ASK1) and translocation of death-associated protein, Daxx, in substantia nigra pars compacta in

- a mouse model of Parkinson's disease: protection by alpha-lipoic acid. *FASEB J* 2007 July;21(9):2226-36.
- (237) Singhal NK, Srivastava G, Agrawal S, Jain SK, Singh MP. Melatonin as a neuroprotective agent in the rodent models of Parkinson's disease: is it all set to irrefutable clinical translation? *Mol Neurobiol* 2012 February;45(1):186-99.
- (238) Squadrito GL, Cueto R, Splenser AE et al. Reaction of uric acid with peroxynitrite and implications for the mechanism of neuroprotection by uric acid. *Arch Biochem Biophys* 2000 April 15;376(2):333-7.
- (239) Cipriani S, Chen X, Schwarzschild MA. Urate: a novel biomarker of Parkinson's disease risk, diagnosis and prognosis. *Biomark Med* 2010 October;4(5):701-12.
- (240) Annanmaki T, Pessala-Driver A, Hokkanen L, Murros K. Uric acid associates with cognition in Parkinson's disease. *Parkinsonism Relat Disord* 2008 November;14(7):576-8.
- (241) Wang XJ, Luo WF, Wang LJ, Mao CJ, Wang L, Liu CF. [Study on uric acid and the related factors associated with cognition in the patients with Parkinson's disease]. *Zhonghua Yi Xue Za Zhi* 2009 June 16;89(23):1633-5.
- (242) Spitsin S, Hooper DC, Leist T, Streletz LJ, Mikheeva T, Koprowski H. Inactivation of peroxynitrite in multiple sclerosis patients after oral administration of inosine may suggest possible approaches to therapy of the disease. *Mult Scler* 2001 October;7(5):313-9.
- (243) Toncev G. Therapeutic value of serum uric acid levels increasing in the treatment of multiple sclerosis. *Vojnosanit Pregl* 2006 October;63(10):879-82.
- (244) Markowitz CE, Spitsin S, Zimmerman V et al. The treatment of multiple sclerosis with inosine. *J Altern Complement Med* 2009 June;15(6):619-25.
- (245) Ikeda Y, Tsuji S, Satoh A, Ishikura M, Shirasawa T, Shimizu T. Protective effects of astaxanthin on 6-hydroxydopamine-induced apoptosis in human neuroblastoma SH-SY5Y cells. *J Neurochem* 2008 December;107(6):1730-40.
- (246) Liu X, Shibata T, Hisaka S, Osawa T. Astaxanthin inhibits reactive oxygen species-mediated cellular toxicity in dopaminergic SH-SY5Y cells via mitochondria-targeted protective mechanism. *Brain Res* 2009 February 13;1254:18-27.
- (247) Liu X, Osawa T. Astaxanthin protects neuronal cells against oxidative damage and is a potent candidate for brain food. *Forum Nutr* 2009;61:129-35.
- (248) Nagesh BG, Kumar A, Singh RL. Chronic pretreatment with acetyl-L-carnitine and +/-DL-alpha-lipoic acid protects against acute glutamate-induced neurotoxicity in rat brain by altering mitochondrial function. *Neurotox Res* 2011 February;19(2):319-29.
- (249) Palacios HH, Yendluri BB, Parvathaneni K et al. Mitochondrion-specific antioxidants as drug treatments for Alzheimer disease. *CNS Neurol Disord Drug Targets* 2011 March;10(2):149-62.

- (250) Liu J, Head E, Gharib AM et al. Memory loss in old rats is associated with brain mitochondrial decay and RNA/DNA oxidation: partial reversal by feeding acetyl-L-carnitine and/or R-alpha - lipoic acid. *Proc Natl Acad Sci U S A* 2002 February 19;99(4):2356-61.
- (251) Ames BN, Liu J. Delaying the mitochondrial decay of aging with acetylcarnitine. *Ann N Y Acad Sci* 2004 November;1033:108-16.
- (252) Aliev G, Liu J, Shenk JC et al. Neuronal mitochondrial amelioration by feeding acetyl-L-carnitine and lipoic acid to aged rats. *J Cell Mol Med* 2009 February;13(2):320-33.
- (253) Long J, Gao F, Tong L, Cotman CW, Ames BN, Liu J. Mitochondrial decay in the brains of old rats: ameliorating effect of alpha-lipoic acid and acetyl-L-carnitine. *Neurochem Res* 2009 April;34(4):755-63.
- (254) Imperato A, Ramacci MT, Angelucci L. Acetyl-L-carnitine enhances acetylcholine release in the striatum and hippocampus of awake freely moving rats. *Neurosci Lett* 1989 December 15;107(1-3):251-5.
- (255) White HL, Scates PW. Acetyl-L-carnitine as a precursor of acetylcholine. *Neurochem Res* 1990 June;15(6):597-601.
- (256) Wawrzencyk A, Nalecz KA, Nalecz MJ. Effect of externally added carnitine on the synthesis of acetylcholine in rat cerebral cortex cells. *Neurochem Int* 1995 June;26(6):635-41.
- (257) Montgomery SA, Thal LJ, Amrein R. Meta-analysis of double blind randomized controlled clinical trials of acetyl-L-carnitine versus placebo in the treatment of mild cognitive impairment and mild Alzheimer's disease. *Int Clin Psychopharmacol* 2003 March;18(2):61-71.
- (258) Levinskaya N, Trenkner E, El IA. Increased GAD-positive neurons in the cortex of taurine-fed mice. *Adv Exp Med Biol* 2006;583:411-7.
- (259) El IA. Taurine improves learning and retention in aged mice. *Neurosci Lett* 2008 May 2;436(1):19-22.
- (260) Saavedra JM. Angiotensin II AT(1) receptor blockers ameliorate inflammatory stress: a beneficial effect for the treatment of brain disorders. *Cell Mol Neurobiol* 2012 July;32(5):667-81.
- (261) Takeda S, Sato N, Takeuchi D et al. Angiotensin receptor blocker prevented beta-amyloid-induced cognitive impairment associated with recovery of neurovascular coupling. *Hypertension* 2009 December;54(6):1345-52.
- (262) Danielyan L, Klein R, Hanson LR et al. Protective effects of intranasal losartan in the APP/PS1 transgenic mouse model of Alzheimer disease. *Rejuvenation Res* 2010 April;13(2-3):195-201.
- (263) Dong YF, Kataoka K, Tokutomi Y et al. Perindopril, a centrally active angiotensin-converting enzyme inhibitor, prevents cognitive impairment in mouse models of Alzheimer's disease. *FASEB J* 2011 September;25(9):2911-20.
- (264) Zhu D, Shi J, Zhang Y et al. Central angiotensin II stimulation promotes beta amyloid production in Sprague Dawley rats. *PLoS ONE* 2011;6(1):e16037.

- (265) Yamada K, Uchida S, Takahashi S et al. Effect of a centrally active angiotensin-converting enzyme inhibitor, perindopril, on cognitive performance in a mouse model of Alzheimer's disease. *Brain Res* 2010 September 17;1352:176-86.
- (266) Kehoe PG, Wilcock GK. Is inhibition of the renin-angiotensin system a new treatment option for Alzheimer's disease? *Lancet Neurol* 2007 April;6(4):373-8.
- (267) Takeda S, Sato N, Ogihara T, Morishita R. The renin-angiotensin system, hypertension and cognitive dysfunction in Alzheimer's disease: new therapeutic potential. *Front Biosci* 2008;13:2253-65.
- (268) Opie LH. Inhibition of the cerebral renin-angiotensin system to limit cognitive decline in elderly hypertensive persons. *Cardiovasc Drugs Ther* 2011 August;25(4):277-9.
- (269) Sink KM, Leng X, Williamson J et al. Angiotensin-converting enzyme inhibitors and cognitive decline in older adults with hypertension: results from the Cardiovascular Health Study. *Arch Intern Med* 2009 July 13;169(13):1195-202.
- (270) Davies NM, Kehoe PG, Ben-Shlomo Y, Martin RM. Associations of anti-hypertensive treatments with Alzheimer's disease, vascular dementia, and other dementias. *J Alzheimers Dis* 2011;26(4):699-708.
- (271) Pelisch N, Hosomi N, Ueno M et al. Blockade of AT1 receptors protects the blood-brain barrier and improves cognition in Dahl salt-sensitive hypertensive rats. *Am J Hypertens* 2011 March;24(3):362-8.
- (272) Lau FC, Bielinski DF, Joseph JA. Inhibitory effects of blueberry extract on the production of inflammatory mediators in lipopolysaccharide-activated BV2 microglia. *J Neurosci Res* 2007 April;85(5):1010-7.
- (273) Poulouse SM, Fisher DR, Larson J et al. Anthocyanin-rich acai (*Euterpe oleracea* Mart.) fruit pulp fractions attenuate inflammatory stress signaling in mouse brain BV-2 microglial cells. *J Agric Food Chem* 2012 February 1;60(4):1084-93.
- (274) Li R, Huang YG, Fang D, Le WD. (-)-Epigallocatechin gallate inhibits lipopolysaccharide-induced microglial activation and protects against inflammation-mediated dopaminergic neuronal injury. *J Neurosci Res* 2004 December 1;78(5):723-31.
- (275) Zheng LT, Ock J, Kwon BM, Suk K. Suppressive effects of flavonoid fisetin on lipopolysaccharide-induced microglial activation and neurotoxicity. *Int Immunopharmacol* 2008 March;8(3):484-94.
- (276) Zhu LH, Bi W, Qi RB, Wang HD, Lu DX. Luteolin inhibits microglial inflammation and improves neuron survival against inflammation. *Int J Neurosci* 2011 June;121(6):329-36.
- (277) Jang S, Dilger RN, Johnson RW. Luteolin inhibits microglia and alters hippocampal-dependent spatial working memory in aged mice. *J Nutr* 2010 October;140(10):1892-8.
- (278) Meng XL, Yang JY, Chen GL et al. Effects of resveratrol and its derivatives on lipopolysaccharide-induced microglial activation and their structure-activity relationships. *Chem Biol Interact* 2008 July 10;174(1):51-9.

- (279) Jang S, Johnson RW. Can consuming flavonoids restore old microglia to their youthful state? *Nutr Rev* 2010 December;68(12):719-28.
- (280) Unno K, Takabayashi F, Yoshida H et al. Daily consumption of green tea catechin delays memory regression in aged mice. *Biogerontology* 2007 April;8(2):89-95.
- (281) Unno K, Ishikawa Y, Takabayashi F et al. Daily ingestion of green tea catechins from adulthood suppressed brain dysfunction in aged mice. *Biofactors* 2008;34(4):263-71.
- (282) Assuncao M, Santos-Marques MJ, Carvalho F, Andrade JP. Green tea averts age-dependent decline of hippocampal signaling systems related to antioxidant defenses and survival. *Free Radic Biol Med* 2010 March 15;48(6):831-8.
- (283) Joseph JA, Shukitt-Hale B, Denisova NA et al. Reversals of age-related declines in neuronal signal transduction, cognitive, and motor behavioral deficits with blueberry, spinach, or strawberry dietary supplementation. *J Neurosci* 1999 September 15;19(18):8114-21.
- (284) Lee JW, Lee YK, Ban JO et al. Green tea (-)-epigallocatechin-3-gallate inhibits beta-amyloid-induced cognitive dysfunction through modification of secretase activity via inhibition of ERK and NF-kappaB pathways in mice. *J Nutr* 2009 October;139(10):1987-93.
- (285) Rezai-Zadeh K, Arendash GW, Hou H et al. Green tea epigallocatechin-3-gallate (EGCG) reduces beta-amyloid mediated cognitive impairment and modulates tau pathology in Alzheimer transgenic mice. *Brain Res* 2008 June 12;1214:177-87.
- (286) Joseph JA, Denisova NA, Arendash G et al. Blueberry supplementation enhances signaling and prevents behavioral deficits in an Alzheimer disease model. *Nutr Neurosci* 2003 June;6(3):153-62.
- (287) Shih PH, Chan YC, Liao JW, Wang MF, Yen GC. Antioxidant and cognitive promotion effects of anthocyanin-rich mulberry (*Morus atropurpurea* L.) on senescence-accelerated mice and prevention of Alzheimer's disease. *J Nutr Biochem* 2010 July;21(7):598-605.
- (288) Wang YJ, Thomas P, Zhong JH et al. Consumption of grape seed extract prevents amyloid-beta deposition and attenuates inflammation in brain of an Alzheimer's disease mouse. *Neurotox Res* 2009 January;15(1):3-14.
- (289) Mori T, Rezai-Zadeh K, Koyama N et al. Tannic acid is a natural beta-secretase inhibitor that prevents cognitive impairment and mitigates Alzheimer-like pathology in transgenic mice. *J Biol Chem* 2012 February 24;287(9):6912-27.
- (290) Hartman RE, Shah A, Fagan AM et al. Pomegranate juice decreases amyloid load and improves behavior in a mouse model of Alzheimer's disease. *Neurobiol Dis* 2006 December;24(3):506-15.
- (291) Chang J, Rimando A, Pallas M et al. Low-dose pterostilbene, but not resveratrol, is a potent neuromodulator in aging and Alzheimer's disease. *Neurobiol Aging* 2012 September;33(9):2062-71.
- (292) Lin CL, Chen TF, Chiu MJ, Way TD, Lin JK. Epigallocatechin gallate (EGCG) suppresses beta-amyloid-induced neurotoxicity through inhibiting c-Abl/FE65 nuclear translocation and GSK3 beta activation. *Neurobiol Aging* 2009 January;30(1):81-92.

- (293) He Y, Cui J, Lee JC et al. Prolonged exposure of cortical neurons to oligomeric amyloid-beta impairs NMDA receptor function via NADPH oxidase-mediated ROS production: protective effect of green tea (-)-epigallocatechin-3-gallate. *ASN Neuro* 2011;3(1):e00050.
- (294) Ansari MA, Abdul HM, Joshi G, Opii WO, Butterfield DA. Protective effect of quercetin in primary neurons against Abeta(1-42): relevance to Alzheimer's disease. *J Nutr Biochem* 2009 April;20(4):269-75.
- (295) Tarozzi A, Merlicco A, Morroni F et al. Cyanidin 3-O-glucopyranoside protects and rescues SH-SY5Y cells against amyloid-beta peptide-induced toxicity. *Neuroreport* 2008 October 8;19(15):1483-6.
- (296) Tarozzi A, Morroni F, Merlicco A et al. Neuroprotective effects of cyanidin 3-O-glucopyranoside on amyloid beta (25-35) oligomer-induced toxicity. *Neurosci Lett* 2010 April 5;473(2):72-6.
- (297) Brewer GJ, Torricelli JR, Lindsey AL et al. Age-related toxicity of amyloid-beta associated with increased pERK and pCREB in primary hippocampal neurons: reversal by blueberry extract. *J Nutr Biochem* 2010 October;21(10):991-8.
- (298) Maher P, Akaishi T, Abe K. Flavonoid fisetin promotes ERK-dependent long-term potentiation and enhances memory. *Proc Natl Acad Sci U S A* 2006 October 31;103(44):16568-73.
- (299) van PH, Lucero MJ, Yeo GW et al. Plant-derived flavanol (-)-epicatechin enhances angiogenesis and retention of spatial memory in mice. *J Neurosci* 2007 May 30;27(22):5869-78.
- (300) Rendeiro C, Vauzour D, Kean RJ et al. Blueberry supplementation induces spatial memory improvements and region-specific regulation of hippocampal BDNF mRNA expression in young rats. *Psychopharmacology (Berl)* 2012 May 9.
- (301) Gotz J, Eckert A, Matamales M, Ittner LM, Liu X. Modes of Abeta toxicity in Alzheimer's disease. *Cell Mol Life Sci* 2011 October;68(20):3359-75.
- (302) Levites Y, Amit T, Mandel S, Youdim MB. Neuroprotection and neurorescue against Abeta toxicity and PKC-dependent release of nonamyloidogenic soluble precursor protein by green tea polyphenol (-)-epigallocatechin-3-gallate. *FASEB J* 2003 May;17(8):952-4.
- (303) Rezai-Zadeh K, Shytle D, Sun N et al. Green tea epigallocatechin-3-gallate (EGCG) modulates amyloid precursor protein cleavage and reduces cerebral amyloidosis in Alzheimer transgenic mice. *J Neurosci* 2005 September 21;25(38):8807-14.
- (304) Obregon DF, Rezai-Zadeh K, Bai Y et al. ADAM10 activation is required for green tea (-)-epigallocatechin-3-gallate-induced alpha-secretase cleavage of amyloid precursor protein. *J Biol Chem* 2006 June 16;281(24):16419-27.
- (305) Fernandez JW, Rezai-Zadeh K, Obregon D, Tan J. EGCG functions through estrogen receptor-mediated activation of ADAM10 in the promotion of non-amyloidogenic processing of APP. *FEBS Lett* 2010 October 8;584(19):4259-67.
- (306) Smith A, Giunta B, Bickford PC, Fountain M, Tan J, Shytle RD. Nanolipidic particles improve the bioavailability and alpha-secretase inducing ability of epigallocatechin-3-gallate (EGCG) for the treatment of Alzheimer's disease. *Int J Pharm* 2010 April 15;389(1-2):207-12.

- (307) Kwak HM, Jeon SY, Sohng BH et al. beta-Secretase (BACE1) inhibitors from pomegranate (*Punica granatum*) husk. *Arch Pharm Res* 2005 December;28(12):1328-32.
- (308) Lee HJ, Seong YH, Bae KH et al. Beta-secretase (BACE1) inhibitors from *Sanguisorbae Radix*. *Arch Pharm Res* 2005 July;28(7):799-803.
- (309) Heber D. Pomegranate Ellagitannins. 2011.
- (310) Rojanathammanee L, Puig KL, Combs CK. Pomegranate polyphenols and extract inhibit nuclear factor of activated T-cell activity and microglial activation in vitro and in a transgenic mouse model of Alzheimer disease. *J Nutr* 2013 May;143(5):597-605.
- (311) Sagara Y, Vanhnasy J, Maher P. Induction of PC12 cell differentiation by flavonoids is dependent upon extracellular signal-regulated kinase activation. *J Neurochem* 2004 September;90(5):1144-55.
- (312) Maher P. Modulation of multiple pathways involved in the maintenance of neuronal function during aging by fisetin. *Genes Nutr* 2009 December;4(4):297-307.
- (313) Kuriyama S, Hozawa A, Ohmori K et al. Green tea consumption and cognitive function: a cross-sectional study from the Tsurugaya Project 1. *Am J Clin Nutr* 2006 February;83(2):355-61.
- (314) Kuriyama S. The relation between green tea consumption and cardiovascular disease as evidenced by epidemiological studies. *J Nutr* 2008 August;138(8):1548S-53S.
- (315) Tanabe N, Suzuki H, Aizawa Y, Seki N. Consumption of green and roasted teas and the risk of stroke incidence: results from the Tokamachi-Nakasato cohort study in Japan. *Int J Epidemiol* 2008 October;37(5):1030-40.
- (316) Arab L, Liu W, Elashoff D. Green and black tea consumption and risk of stroke: a meta-analysis. *Stroke* 2009 May;40(5):1786-92.
- (317) Liang W, Lee AH, Binns CW, Huang R, Hu D, Zhou Q. Tea consumption and ischemic stroke risk: a case-control study in southern China. *Stroke* 2009 July;40(7):2480-5.
- (318) Suzuki E, Yorifuji T, Takao S et al. Green tea consumption and mortality among Japanese elderly people: the prospective Shizuoka elderly cohort. *Ann Epidemiol* 2009 October;19(10):732-9.
- (319) Tomata Y, Kakizaki M, Nakaya N et al. Green tea consumption and the risk of incident functional disability in elderly Japanese: the Ohsaki Cohort 2006 Study. *Am J Clin Nutr* 2012 March;95(3):732-9.
- (320) Letenneur L, Proust-Lima C, Le GA, Dartigues JF, Barberger-Gateau P. Flavonoid intake and cognitive decline over a 10-year period. *Am J Epidemiol* 2007 June 15;165(12):1364-71.
- (321) Devore EE, Kang JH, Breteler MM, Grodstein F. Dietary intakes of berries and flavonoids in relation to cognitive decline. *Ann Neurol* 2012 July;72(1):135-43.
- (322) Santos AC, Veiga F, Ribeiro AJ. New delivery systems to improve the bioavailability of resveratrol. *Expert Opin Drug Deliv* 2011 August;8(8):973-90.

- (323) Kapetanovic IM, Muzzio M, Huang Z, Thompson TN, McCormick DL. Pharmacokinetics, oral bioavailability, and metabolic profile of resveratrol and its dimethylether analog, pterostilbene, in rats. *Cancer Chemother Pharmacol* 2011 September;68(3):593-601.
- (324) Hopper RA, Garthwaite J. Tonic and phasic nitric oxide signals in hippocampal long-term potentiation. *J Neurosci* 2006 November 8;26(45):11513-21.
- (325) Brown GC, Neher JJ. Inflammatory neurodegeneration and mechanisms of microglial killing of neurons. *Mol Neurobiol* 2010 June;41(2-3):242-7.
- (326) Austin SA, Santhanam AV, Katusic ZS. Endothelial nitric oxide modulates expression and processing of amyloid precursor protein. *Circ Res* 2010 December 10;107(12):1498-502.
- (327) McCarty MF. Vascular nitric oxide may lessen Alzheimer's risk. *Med Hypotheses* 1998 December;51(6):465-76.
- (328) Taqatqeh F, Mergia E, Neitz A, Eysel UT, Koesling D, Mittmann T. More than a retrograde messenger: nitric oxide needs two cGMP pathways to induce hippocampal long-term potentiation. *J Neurosci* 2009 July 22;29(29):9344-50.
- (329) Boran MS, Baltrons MA, Garcia A. The ANP-cGMP-protein kinase G pathway induces a phagocytic phenotype but decreases inflammatory gene expression in microglial cells. *Glia* 2008 March;56(4):394-411.
- (330) Yoshioka Y, Takeda N, Yamamuro A, Kasai A, Maeda S. Nitric oxide inhibits lipopolysaccharide-induced inducible nitric oxide synthase expression and its own production through the cGMP signaling pathway in murine microglia BV-2 cells. *J Pharmacol Sci* 2010;113(2):153-60.
- (331) Xin ZC, Kim EK, Lin CS et al. Effects of icariin on cGMP-specific PDE5 and cAMP-specific PDE4 activities. *Asian J Androl* 2003 March;5(1):15-8.
- (332) Jiang Z, Hu B, Wang J et al. Effect of icariin on cyclic GMP levels and on the mRNA expression of cGMP-binding cGMP-specific phosphodiesterase (PDE5) in penile cavernosum. *J Huazhong Univ Sci Technolog Med Sci* 2006;26(4):460-2.
- (333) Zhang J, Wang YB, Ma CG et al. Icarisid II, a PDE5 inhibitor from *Epimedium wanshanense*, increases cellular cGMP by enhancing NOS in diabetic ED rats corpus cavernosum tissue. *Andrologia* 2012 May;44 Suppl 1:87-93.
- (334) Puzzo D, Vitolo O, Trinchese F, Jacob JP, Palmeri A, Arancio O. Amyloid-beta peptide inhibits activation of the nitric oxide/cGMP/cAMP-responsive element-binding protein pathway during hippocampal synaptic plasticity. *J Neurosci* 2005 July 20;25(29):6887-97.
- (335) Monfort P, Felipe V. Amyloid-beta impairs, and ibuprofen restores, the cGMP pathway, synaptic expression of AMPA receptors and long-term potentiation in the hippocampus. *J Alzheimers Dis* 2010;22(3):795-809.
- (336) Miller TW, Isenberg JS, Shih HB, Wang Y, Roberts DD. Amyloid-beta inhibits No-cGMP signaling in a. *PLoS ONE* 2010;5(12):e15686.

- (337) Puzzo D, Staniszewski A, Deng SX et al. Phosphodiesterase 5 inhibition improves synaptic function, memory, and amyloid-beta load in an Alzheimer's disease mouse model. *J Neurosci* 2009 June 24;29(25):8075-86.
- (338) Cuadrado-Tejedor M, Hervias I, Ricobaraza A et al. Sildenafil restores cognitive function without affecting beta-amyloid burden in a mouse model of Alzheimer's disease. *Br J Pharmacol* 2011 December;164(8):2029-41.
- (339) Rutten K, Vente JD, Sik A, Ittersum MM, Prickaerts J, Blokland A. The selective PDE5 inhibitor, sildenafil, improves object memory in Swiss mice and increases cGMP levels in hippocampal slices. *Behav Brain Res* 2005 October 14;164(1):11-6.
- (340) Nie J, Luo Y, Huang XN, Gong QH, Wu Q, Shi JS. Icariin inhibits beta-amyloid peptide segment 25-35 induced expression of beta-secretase in rat hippocampus. *Eur J Pharmacol* 2010 January 25;626(2-3):213-8.
- (341) Urano T, Tohda C. Icariin improves memory impairment in Alzheimer's disease model mice (5xFAD) and attenuates amyloid beta-induced neurite atrophy. *Phytother Res* 2010 November;24(11):1658-63.
- (342) He XL, Zhou WQ, Bi MG, Du GH. Neuroprotective effects of icariin on memory impairment and neurochemical deficits in senescence-accelerated mouse prone 8 (SAMP8) mice. *Brain Res* 2010 June 2;1334:73-83.
- (343) Sabayan B, Zamiri N, Farshchizarabi S, Sabayan B. Phosphodiesterase-5 inhibitors: novel weapons against Alzheimer's disease? *Int J Neurosci* 2010 December;120(12):746-51.
- (344) Zhao S, Zhang L, Lian G et al. Sildenafil attenuates LPS-induced pro-inflammatory responses through down-regulation of intracellular ROS-related MAPK/NF-kappaB signaling pathways in N9 microglia. *Int Immunopharmacol* 2011 April;11(4):468-74.
- (345) Zeng KW, Fu H, Liu GX, Wang XM. Icariin attenuates lipopolysaccharide-induced microglial activation and resultant death of neurons by inhibiting TAK1/IKK/NF-kappaB and JNK/p38 MAPK pathways. *Int Immunopharmacol* 2010 June;10(6):668-78.
- (346) Forstermann U. Janus-faced role of endothelial NO synthase in vascular disease: uncoupling of oxygen reduction from NO synthesis and its pharmacological reversal. *Biol Chem* 2006 December;387(12):1521-33.
- (347) Hollenberg K. Vascular action of cocoa flavanols in humans: the roots of the story. *J Cardiovasc Pharmacol* 2006;47 Suppl 2:S99-102.
- (348) Hollenberg NK, Fisher ND, McCullough ML. Flavanols, the Kuna, cocoa consumption, and nitric oxide. *J Am Soc Hypertens* 2009 March;3(2):105-12.
- (349) Schroeter H, Heiss C, Balzer J et al. (-)-Epicatechin mediates beneficial effects of flavanol-rich cocoa on vascular function in humans. *Proc Natl Acad Sci U S A* 2006 January 24;103(4):1024-9.
- (350) Loke WM, Hodgson JM, Proudfoot JM, McKinley AJ, Puddey IB, Croft KD. Pure dietary flavonoids quercetin and (-)-epicatechin augment nitric oxide products and reduce endothelin-1 acutely in healthy men. *Am J Clin Nutr* 2008 October;88(4):1018-25.

- (351) Oyama J, Maeda T, Kouzuma K et al. Green tea catechins improve human forearm endothelial dysfunction and have antiatherosclerotic effects in smokers. *Circ J* 2010 March;74(3):578-88.
- (352) Park CS, Kim W, Woo JS et al. Green tea consumption improves endothelial function but not circulating endothelial progenitor cells in patients with chronic renal failure. *Int J Cardiol* 2010 November 19;145(2):261-2.
- (353) Alexopoulos N, Vlachopoulos C, Aznaouridis K et al. The acute effect of green tea consumption on endothelial function in healthy individuals. *Eur J Cardiovasc Prev Rehabil* 2008 June;15(3):300-5.
- (354) Levenson JM, O'Riordan KJ, Brown KD, Trinh MA, Molfese DL, Sweatt JD. Regulation of histone acetylation during memory formation in the hippocampus. *J Biol Chem* 2004 September 24;279(39):40545-59.
- (355) Fischer A, Sananbenesi F, Wang X, Dobbin M, Tsai LH. Recovery of learning and memory is associated with chromatin remodelling. *Nature* 2007 May 10;447(7141):178-82.
- (356) Guan JS, Haggarty SJ, Giacometti E et al. HDAC2 negatively regulates memory formation and synaptic plasticity. *Nature* 2009 May 7;459(7243):55-60.
- (357) Graff J, Rei D, Guan JS et al. An epigenetic blockade of cognitive functions in the neurodegenerating brain. *Nature* 2012 March 8;483(7388):222-6.
- (358) Xu K, Dai XL, Huang HC, Jiang ZF. Targeting HDACs: a promising therapy for Alzheimer's disease. *Oxid Med Cell Longev* 2011;2011:143269.
- (359) Kilgore M, Miller CA, Fass DM et al. Inhibitors of class 1 histone deacetylases reverse contextual memory deficits in a mouse model of Alzheimer's disease. *Neuropsychopharmacology* 2010 March;35(4):870-80.
- (360) Govindarajan N, Agis-Balboa RC, Walter J, Sananbenesi F, Fischer A. Sodium butyrate improves memory function in an Alzheimer's disease mouse model when administered at an advanced stage of disease progression. *J Alzheimers Dis* 2011;26(1):187-97.
- (361) Wiley JC, Pettan-Brewer C, Ladiges WC. Phenylbutyric acid reduces amyloid plaques and rescues cognitive behavior in AD transgenic mice. *Aging Cell* 2011 June;10(3):418-28.
- (362) Ricobaraza A, Cuadrado-Tejedor M, Marco S, Perez-Otano I, Garcia-Osta A. Phenylbutyrate rescues dendritic spine loss associated with memory deficits in a mouse model of Alzheimer disease. *Hippocampus* 2012 May;22(5):1040-50.
- (363) Ricobaraza A, Cuadrado-Tejedor M, Perez-Mediavilla A, Frechilla D, Del RJ, Garcia-Osta A. Phenylbutyrate ameliorates cognitive deficit and reduces tau pathology in an Alzheimer's disease mouse model. *Neuropsychopharmacology* 2009 June;34(7):1721-32.
- (364) Fontan-Lozano A, Romero-Granados R, Troncoso J, Munera A, Delgado-Garcia JM, Carrion AM. Histone deacetylase inhibitors improve learning consolidation in young and in KA-induced-neurodegeneration and SAMP-8-mutant mice. *Mol Cell Neurosci* 2008 October;39(2):193-201.

- (365) Zeng Y, Tan M, Kohyama J et al. Epigenetic enhancement of BDNF signaling rescues synaptic plasticity in aging. *J Neurosci* 2011 December 7;31(49):17800-10.
- (366) Egorin MJ, Yuan ZM, Sentz DL, Plaisance K, Eiseman JL. Plasma pharmacokinetics of butyrate after intravenous administration of sodium butyrate or oral administration of tributyrin or sodium butyrate to mice and rats. *Cancer Chemother Pharmacol* 1999;43(6):445-53.
- (367) Edelman MJ, Bauer K, Khanwani S et al. Clinical and pharmacologic study of tributyrin: an oral butyrate prodrug. *Cancer Chemother Pharmacol* 2003 May;51(5):439-44.
- (368) Su J, He L, ZHANG N, Ho PC. Evaluation of tributyrin lipid emulsion with affinity to low-density lipoprotein: pharmacokinetics in adult male Wistar rats and cellular activity on Caco-2 and HepG2 cell lines. *J Pharmacol Exp Ther* 2006 January;316(1):62-70.
- (369) Vinolo MA, Rodrigues H, Festuccia WT et al. Tributyrin attenuates obesity-associated inflammation and insulin resistance in high-fat fed mice. *Am J Physiol Endocrinol Metab* 2012 May 22.
- (370) Hogarth P, Lovrecic L, Krainc D. Sodium phenylbutyrate in Huntington's disease: a dose-finding study. *Mov Disord* 2007 October 15;22(13):1962-4.
- (371) Cudkowicz ME, Andres PL, Macdonald SA et al. Phase 2 study of sodium phenylbutyrate in ALS. *Amyotroph Lateral Scler* 2009 April;10(2):99-106.
- (372) Pomponi MG, Neri G. Butyrate and acetyl-carnitine inhibit the cytogenetic expression of the fragile X in vitro. *Am J Med Genet* 1994 July 15;51(4):447-50.
- (373) Madiraju P, Pande SV, Prentki M, Madiraju SR. Mitochondrial acetylcarnitine provides acetyl groups for nuclear histone acetylation. *Epigenetics* 2009 August 16;4(6):399-403.
- (374) Chiechio S, Copani A, Zammataro M, Battaglia G, Gereau RW, Nicoletti F. Transcriptional regulation of type-2 metabotropic glutamate receptors: an epigenetic path to novel treatments for chronic pain. *Trends Pharmacol Sci* 2010 April;31(4):153-60.
- (375) Nasca C, Xenos D, Barone Y et al. L-acetylcarnitine causes rapid antidepressant effects through the epigenetic induction of mGlu2 receptors. *Proc Natl Acad Sci U S A* 2013 March 19;110(12):4804-9.
- (376) Rapoport M, Dawson HN, Binder LI, Vitek MP, Ferreira A. Tau is essential to beta -amyloid-induced neurotoxicity. *Proc Natl Acad Sci U S A* 2002 April 30;99(9):6364-9.
- (377) Roberson ED, Scarce-Levie K, Palop JJ et al. Reducing endogenous tau ameliorates amyloid beta-induced deficits in an Alzheimer's disease mouse model. *Science* 2007 May 4;316(5825):750-4.
- (378) Roberson ED, Halabisky B, Yoo JW et al. Amyloid-beta/Fyn-induced synaptic, network, and cognitive impairments depend on tau levels in multiple mouse models of Alzheimer's disease. *J Neurosci* 2011 January 12;31(2):700-11.
- (379) Hong M, Chen DC, Klein PS, Lee VM. Lithium reduces tau phosphorylation by inhibition of glycogen synthase kinase-3. *J Biol Chem* 1997 October 3;272(40):25326-32.

- (380) Rockenstein E, Torrance M, Adame A et al. Neuroprotective effects of regulators of the glycogen synthase kinase-3beta signaling pathway in a transgenic model of Alzheimer's disease are associated with reduced amyloid precursor protein phosphorylation. *J Neurosci* 2007 February 21;27(8):1981-91.
- (381) Medina M, Avila J. Glycogen synthase kinase-3 (GSK-3) inhibitors for the treatment of Alzheimer's disease. *Curr Pharm Des* 2010;16(25):2790-8.
- (382) Nunes PV, Forlenza OV, Gattaz WF. Lithium and risk for Alzheimer's disease in elderly patients with bipolar disorder. *Br J Psychiatry* 2007 April;190:359-60.
- (383) Kessing LV, Sondergard L, Forman JL, Andersen PK. Lithium treatment and risk of dementia. *Arch Gen Psychiatry* 2008 November;65(11):1331-5.
- (384) Kessing LV, Forman JL, Andersen PK. Does lithium protect against dementia? *Bipolar Disord* 2010 February;12(1):87-94.
- (385) Forlenza OV, Diniz BS, Radanovic M, Santos FS, Talib LL, Gattaz WF. Disease-modifying properties of long-term lithium treatment for amnesic mild cognitive impairment: randomised controlled trial. *Br J Psychiatry* 2011 May;198(5):351-6.
- (386) Caccamo A, Oddo S, Tran LX, Laferla FM. Lithium reduces tau phosphorylation but not A beta or working memory deficits in a transgenic model with both plaques and tangles. *Am J Pathol* 2007 May;170(5):1669-75.
- (387) Sudduth TL, Wilson JG, Everhart A, Colton CA, Wilcock DM. Lithium treatment of APPSwDI/NOS2-/- mice leads to reduced hyperphosphorylated tau, increased amyloid deposition and altered inflammatory phenotype. *PLoS ONE* 2012;7(2):e31993.
- (388) Tajés M, Gutierrez-Cuesta J, Folch J et al. Lithium treatment decreases activities of tau kinases in a murine model of senescence. *J Neuropathol Exp Neurol* 2008 June;67(6):612-23.
- (389) Zhang X, Heng X, Li T et al. Long-term treatment with lithium alleviates memory deficits and reduces amyloid-beta production in an aged Alzheimer's disease transgenic mouse model. *J Alzheimers Dis* 2011;24(4):739-49.
- (390) Nunes MA, Viel TA, Buck HS. Microdose lithium treatment stabilized cognitive impairment in patients with Alzheimers disease. *Curr Alzheimer Res* 2012 June 29.
- (391) Corcoran NM, Martin D, Hutter-Paier B et al. Sodium selenate specifically activates PP2A phosphatase, dephosphorylates tau and reverses memory deficits in an Alzheimer's disease model. *J Clin Neurosci* 2010 August;17(8):1025-33.
- (392) van EJ, Ke YD, Liu X et al. Sodium selenate mitigates tau pathology, neurodegeneration, and functional deficits in Alzheimer's disease models. *Proc Natl Acad Sci U S A* 2010 August 3;107(31):13888-93.
- (393) Corcoran NM, Hovens CM, Michael M, Rosenthal MA, Costello AJ. Open-label, phase I dose-escalation study of sodium selenate, a novel activator of PP2A, in patients with castration-resistant prostate cancer. *Br J Cancer* 2010 August 10;103(4):462-8.

- (394) McGeer PL, Schulzer M, McGeer EG. Arthritis and anti-inflammatory agents as possible protective factors for Alzheimer's disease: a review of 17 epidemiologic studies. *Neurology* 1996 August;47(2):425-32.
- (395) Etminan M, Gill S, Samii A. Effect of non-steroidal anti-inflammatory drugs on risk of Alzheimer's disease: systematic review and meta-analysis of observational studies. *BMJ* 2003 July 19;327(7407):128.
- (396) Szekely CA, Thorne JE, Zandi PP et al. Nonsteroidal anti-inflammatory drugs for the prevention of Alzheimer's disease: a systematic review. *Neuroepidemiology* 2004 July;23(4):159-69.
- (397) Jaturapatporn D, Isaac MG, McCleery J, Tabet N. Aspirin, steroidal and non-steroidal anti-inflammatory drugs for the treatment of Alzheimer's disease. *Cochrane Database Syst Rev* 2012;2:CD006378.
- (398) Ikonovic MD, Abrahamson EE, Uz T, Manev H, DeKosky ST. Increased 5-lipoxygenase immunoreactivity in the hippocampus of patients with Alzheimer's disease. *J Histochem Cytochem* 2008 December;56(12):1065-73.
- (399) Puccio S, Chu J, Pratico D. Involvement of 5-lipoxygenase in the corticosteroid-dependent amyloid beta formation: in vitro and in vivo evidence. *PLoS ONE* 2011;6(1):e15163.
- (400) Chu J, Giannopoulos PF, Ceballos-Diaz C, Golde TE, Pratico D. Adeno-associated virus-mediated brain delivery of 5-lipoxygenase modulates the AD-like phenotype of APP mice. *Mol Neurodegener* 2012;7(1):1.
- (401) Wang ZJ, Zhou B, Mao WW, Yin M. Overexpression of 5-lipoxygenase increases the neuronal vulnerability of PC12 cells to Aβ(4)(2). *Yakugaku Zasshi* 2011;131(12):1843-53.
- (402) Firuzi O, Zhuo J, Chinnici CM, Wisniewski T, Pratico D. 5-Lipoxygenase gene disruption reduces amyloid-beta pathology in a mouse model of Alzheimer's disease. *FASEB J* 2008 April;22(4):1169-78.
- (403) Chu J, Pratico D. 5-lipoxygenase as an endogenous modulator of amyloid beta formation in vivo. *Ann Neurol* 2011 January;69(1):34-46.
- (404) Chu J, Pratico D. Pharmacologic blockade of 5-lipoxygenase improves the amyloidotic phenotype of an Alzheimer's disease transgenic mouse model involvement of gamma-secretase. *Am J Pathol* 2011 April;178(4):1762-9.
- (405) Chu J, Pratico D. Involvement of 5-lipoxygenase activating protein in the amyloidotic phenotype of an Alzheimer's disease mouse model. *J Neuroinflammation* 2012;9:127.
- (406) Chu J, Pratico D. 5-Lipoxygenase pharmacological blockade decreases tau phosphorylation in vivo: involvement of the cyclin-dependent kinase-5. *Neurobiol Aging* 2013 January 16.
- (407) Berger W, De Chandt MT, Cairns CB. Zileuton: clinical implications of 5-Lipoxygenase inhibition in severe airway disease. *Int J Clin Pract* 2007 April;61(4):663-76.
- (408) Safayhi H, Sailer ER, Ammon HP. 5-Lipoxygenase inhibition by acetyl-11-keto-beta-boswellic acid (AKBA) by a novel mechanism. *Phytomedicine* 1996 May;3(1):71-2.

- (409) Siemoneit U, Pergola C, Jazzar B et al. On the interference of boswellic acids with 5-lipoxygenase: mechanistic studies in vitro and pharmacological relevance. *Eur J Pharmacol* 2009 March 15;606(1-3):246-54.
- (410) Abdel-Tawab M, Werz O, Schubert-Zsilavecz M. Boswellia serrata: an overall assessment of in vitro, preclinical, pharmacokinetic and clinical data. *Clin Pharmacokinet* 2011 June;50(6):349-69.
- (411) Siemoneit U, Pergola C, Jazzar B et al. On the interference of boswellic acids with 5-lipoxygenase: mechanistic studies in vitro and pharmacological relevance. *Eur J Pharmacol* 2009 March 15;606(1-3):246-54.
- (412) Siemoneit U, Koeberle A, Rossi A et al. Inhibition of microsomal prostaglandin E2 synthase-1 as a molecular basis for the anti-inflammatory actions of boswellic acids from frankincense. *Br J Pharmacol* 2011 January;162(1):147-62.
- (413) Bishnoi M, Patil CS, Kumar A, Kulkarni SK. Protective effects of nimesulide (COX Inhibitor), AKBA (5-LOX Inhibitor), and their combination in aging-associated abnormalities in mice. *Methods Find Exp Clin Pharmacol* 2005 September;27(7):465-70.
- (414) Sugaya K, Uz T, Kumar V, Manev H. New anti-inflammatory treatment strategy in Alzheimer's disease. *Jpn J Pharmacol* 2000 February;82(2):85-94.
- (415) Kremer JM, Jubiz W, Michalek A et al. Fish-oil fatty acid supplementation in active rheumatoid arthritis. A double-blinded, controlled, crossover study. *Ann Intern Med* 1987 April;106(4):497-503.
- (416) Cansev M, Watkins CJ, van der Beek EM, Wurtman RJ. Oral uridine-5'-monophosphate (UMP) increases brain CDP-choline levels in gerbils. *Brain Res* 2005 October 5;1058(1-2):101-8.
- (417) Sakamoto T, Cansev M, Wurtman RJ. Oral supplementation with docosahexaenoic acid and uridine-5'-monophosphate increases dendritic spine density in adult gerbil hippocampus. *Brain Res* 2007 November 28;1182:50-9.
- (418) Holguin S, Martinez J, Chow C, Wurtman R. Dietary uridine enhances the improvement in learning and memory produced by administering DHA to gerbils. *FASEB J* 2008 November;22(11):3938-46.
- (419) Wurtman RJ, Cansev M, Sakamoto T, Ulus I. Nutritional modifiers of aging brain function: use of uridine and other phosphatide precursors to increase formation of brain synapses. *Nutr Rev* 2010 December;68 Suppl 2:S88-101.
- (420) Dileepan KN, Kennedy J. Rapid conversion of newly-synthesized orotate to uridine-5-monophosphate by rat liver cytosolic enzymes. *FEBS Lett* 1983 March 7;153(1):1-5.
- (421) Rosenfeldt FL, Richards SM, Lin Z, Pepe S, Conyers RA. Mechanism of cardioprotective effect of orotic acid. *Cardiovasc Drugs Ther* 1998 September;12 Suppl 2:159-70.
- (422) Desi I, Szelenyi I, Sos J. [Influence of magnesium-orotate and orotic acid on learning processes in normal and neurotoxically injured rats]. *Arzneimittelforschung* 1971 June;21(6):780-5.

- (423) Ruthrich HL, Wetzel W, Matthies H. The persistence of the effect of methylglucamine orotate on the retention of a learned behavior in the rat. *Acta Biol Med Ger* 1980;39(4):415-8.
- (424) Ruthrich HL, Wetzel W, Matthies H. Memory retention in old rats: improvement by orotic acid. *Psychopharmacology (Berl)* 1983;79(4):348-51.
- (425) Krug M, Koch M, Schoof E, Wagner M, Matthies H. Methylglucamine orotate, a memory-improving drug, prolongs hippocampal long-term potentiation. *Eur J Pharmacol* 1989 December 7;173(2-3):223-6.
- (426) Rosenfeldt FL. Metabolic supplementation with orotic acid and magnesium orotate. *Cardiovasc Drugs Ther* 1998 September;12 Suppl 2:147-52.
- (427) Geiss KR, Stergiou N, Jester, Neuenfeld HU, Jester HG. Effects of magnesium orotate on exercise tolerance in patients with coronary heart disease. *Cardiovasc Drugs Ther* 1998 September;12 Suppl 2:153-6.
- (428) Stepura OB, Martynow AI. Magnesium orotate in severe congestive heart failure (MACH). *Int J Cardiol* 2009 May 1;134(1):145-7.
- (429) Slutsky I, Abumaria N, Wu LJ et al. Enhancement of learning and memory by elevating brain magnesium. *Neuron* 2010 January 28;65(2):165-77.
- (430) Abumaria N, Yin B, Zhang L et al. Effects of elevation of brain magnesium on fear conditioning, fear extinction, and synaptic plasticity in the infralimbic prefrontal cortex and lateral amygdala. *J Neurosci* 2011 October 19;31(42):14871-81.
- (431) Li W, Yu J, Liu Y et al. Elevation of brain magnesium prevents and reverses cognitive deficits and synaptic loss in Alzheimer's disease mouse model. *J Neurosci* 2013 May 8;33(19):8423-41.
- (432) Yu J, Sun M, Chen Z et al. Magnesium modulates amyloid-beta protein precursor trafficking and processing. *J Alzheimers Dis* 2010;20(4):1091-106.
- (433) Lee M, Jantaratnotai N, McGeer E, McLarnon JG, McGeer PL. Mg²⁺ ions reduce microglial and THP-1 cell neurotoxicity by inhibiting Ca²⁺ entry through purinergic channels. *Brain Res* 2011 January 19;1369:21-35.
- (434) Lemke MR. Plasma magnesium decrease and altered calcium/magnesium ratio in severe dementia of the Alzheimer type. *Biol Psychiatry* 1995 March 1;37(5):341-3.
- (435) Andradi E, Igaz S, Molnar Z, Mako S. Disturbances of magnesium concentrations in various brain areas in Alzheimer's disease. *Magnes Res* 2000 September;13(3):189-96.
- (436) Andradi E, Pali N, Molnar Z, Kosel S. Brain aluminum, magnesium and phosphorus contents of control and Alzheimer-diseased patients. *J Alzheimers Dis* 2005 August;7(4):273-84.
- (437) Parnetti L, Mignini F, Tomassoni D, Traini E, Amenta F. Cholinergic precursors in the treatment of cognitive impairment of vascular origin: ineffective approaches or need for re-evaluation? *J Neurol Sci* 2007 June 15;257(1-2):264-9.

- (438) Fioravanti M, Buckley AE. Citicoline (Cognizin) in the treatment of cognitive impairment. *Clin Interv Aging* 2006;1(3):247-51.
- (439) Parnetti L, Amenta F, Gallai V. Choline alphoscerate in cognitive decline and in acute cerebrovascular disease: an analysis of published clinical data. *Mech Ageing Dev* 2001 November;122(16):2041-55.
- (440) Bai DL, Tang XC, He XC. Huperzine A, a potential therapeutic agent for treatment of Alzheimer's disease. *Curr Med Chem* 2000 March;7(3):355-74.
- (441) Wang BS, Wang H, Wei ZH, Song YY, Zhang L, Chen HZ. Efficacy and safety of natural acetylcholinesterase inhibitor huperzine A in the treatment of Alzheimer's disease: an updated meta-analysis. *J Neural Transm* 2009 April;116(4):457-65.
- (442) Wang R, Tang XC. Neuroprotective effects of huperzine A. A natural cholinesterase inhibitor for the treatment of Alzheimer's disease. *Neurosignals* 2005;14(1-2):71-82.
- (443) Wang CY, Zheng W, Wang T et al. Huperzine A activates Wnt/beta-catenin signaling and enhances the nonamyloidogenic pathway in an Alzheimer transgenic mouse model. *Neuropsychopharmacology* 2011 April;36(5):1073-89.
- (444) Wang Y, Tang XC, Zhang HY. Huperzine A alleviates synaptic deficits and modulates amyloidogenic and nonamyloidogenic pathways in APPswe/PS1dE9 transgenic mice. *J Neurosci Res* 2012 February;90(2):508-17.
- (445) McQuail JA, Riddle DR, Nicolle MM. Neuroinflammation not associated with cholinergic degeneration in aged-impaired brain. *Neurobiol Aging* 2011 December;32(12):2322-4.
- (446) Matos M, Augusto E, Machado NJ, Dos Santos-Rodrigues A, Cunha RA, Agostinho P. Astrocytic Adenosine A2A Receptors Control the Amyloid-beta Peptide-Induced Decrease of Glutamate Uptake. *J Alzheimers Dis* 2012 January 1;31(3):555-67.
- (447) Hellweg R, Wirth Y, Janetzky W, Hartmann S. Efficacy of memantine in delaying clinical worsening in Alzheimer's disease (AD): responder analyses of nine clinical trials with patients with moderate to severe AD. *Int J Geriatr Psychiatry* 2012 June;27(6):651-6.
- (448) Schmidt R, Ropele S, Pendl B et al. Longitudinal multimodal imaging in mild to moderate Alzheimer disease: a pilot study with memantine. *J Neurol Neurosurg Psychiatry* 2008 December;79(12):1312-7.
- (449) Weiner MW, Sadowsky C, Saxton J et al. Magnetic resonance imaging and neuropsychological results from a trial of memantine in Alzheimer's disease. *Alzheimers Dement* 2011 July;7(4):425-35.
- (450) Schneider LS, Dagerman KS, Higgins JP, McShane R. Lack of evidence for the efficacy of memantine in mild Alzheimer disease. *Arch Neurol* 2011 August;68(8):991-8.
- (451) Lipton SA. The molecular basis of memantine action in Alzheimer's disease and other neurologic disorders: low-affinity, uncompetitive antagonism. *Curr Alzheimer Res* 2005 April;2(2):155-65.

- (452) Lee MS, Kwon YT, Li M, Peng J, Friedlander RM, Tsai LH. Neurotoxicity induces cleavage of p35 to p25 by calpain. *Nature* 2000 May 18;405(6784):360-4.
- (453) Granic I, Nyakas C, Luiten PG et al. Calpain inhibition prevents amyloid-beta-induced neurodegeneration and associated behavioral dysfunction in rats. *Neuropharmacology* 2010 September;59(4-5):334-42.
- (454) Nikkel AL, Martino B, Markosyan S et al. The novel calpain inhibitor A-705253 prevents stress-induced tau hyperphosphorylation in vitro and in vivo. *Neuropharmacology* 2012 September;63(4):606-12.
- (455) Medeiros R, Kitazawa M, Chabrier MA et al. Calpain Inhibitor A-705253 Mitigates Alzheimer's Disease-Like Pathology and Cognitive Decline in Aged 3xTgAD Mice. *Am J Pathol* 2012 August;181(2):616-25.
- (456) Jang SW, Liu X, Yepes M et al. A selective TrkB agonist with potent neurotrophic activities by 7,8-dihydroxyflavone. *Proc Natl Acad Sci U S A* 2010 February 9;107(6):2687-92.
- (457) Minichiello L. TrkB signalling pathways in LTP and learning. *Nat Rev Neurosci* 2009 December;10(12):850-60.
- (458) Devi L, Ohno M. 7,8-dihydroxyflavone, a small-molecule TrkB agonist, reverses memory deficits and BACE1 elevation in a mouse model of Alzheimer's disease. *Neuropsychopharmacology* 2012 January;37(2):434-44.
- (459) Zeng Y, Lv F, Li L, Yu H, Dong M, Fu Q. 7,8-dihydroxyflavone rescues spatial memory and synaptic plasticity in cognitively impaired aged rats. *J Neurochem* 2012 August;122(4):800-11.
- (460) Jang SW, Liu X, Chan CB et al. Deoxygedunin, a natural product with potent neurotrophic activity in mice. *PLoS ONE* 2010;5(7):e11528.
- (461) Tohda C, Joyashiki E. Sominone enhances neurite outgrowth and spatial memory mediated by the neurotrophic factor receptor, RET. *Br J Pharmacol* 2009 August;157(8):1427-40.
- (462) Joyashiki E, Matsuya Y, Tohda C. Sominone improves memory impairments and increases axonal density in Alzheimer's disease model mice, 5XFAD. *Int J Neurosci* 2011 April;121(4):181-90.
- (463) Shemesh E, Rudich A, Harman-Boehm I, Cukierman-Yaffe T. Effect of intranasal insulin on cognitive function: a systematic review. *J Clin Endocrinol Metab* 2012 February;97(2):366-76.
- (464) Schioth HB, Craft S, Brooks SJ, Frey WH, Benedict C. Brain Insulin Signaling and Alzheimer's Disease: Current Evidence and Future Directions. *Mol Neurobiol* 2011 December 29.
- (465) Reger MA, Watson GS, Green PS et al. Intranasal insulin improves cognition and modulates beta-amyloid in early AD. *Neurology* 2008 February 5;70(6):440-8.
- (466) Craft S, Baker LD, Montine TJ et al. Intranasal insulin therapy for Alzheimer disease and amnesic mild cognitive impairment: a pilot clinical trial. *Arch Neurol* 2012 January;69(1):29-38.

- (467) Mawuenyega KG, Sigurdson W, Ovod V et al. Decreased clearance of CNS beta-amyloid in Alzheimer's disease. *Science* 2010 December 24;330(6012):1774.
- (468) Deane R, Bell RD, Sagare A, Zlokovic BV. Clearance of amyloid-beta peptide across the blood-brain barrier: implication for therapies in Alzheimer's disease. *CNS Neurol Disord Drug Targets* 2009 March;8(1):16-30.
- (469) Sagare AP, Deane R, Zlokovic BV. Low-density lipoprotein receptor-related protein 1: A physiological Abeta homeostatic mechanism with multiple therapeutic opportunities. *Pharmacol Ther* 2012 July 20.
- (470) Sehgal N, Gupta A, Valli RK et al. Withania somnifera reverses Alzheimer's disease pathology by enhancing low-density lipoprotein receptor-related protein in liver. *Proc Natl Acad Sci U S A* 2012 February 28;109(9):3510-5.
- (471) Qosa H, Abuznait AH, Hill RA, Kaddoumi A. Enhanced Brain Amyloid-beta Clearance by Rifampicin and Caffeine as a Possible Protective Mechanism Against Alzheimer's Disease. *J Alzheimers Dis* 2012 January 1;31(1):151-65.
- (472) Loeb MB, Molloy DW, Smieja M et al. A randomized, controlled trial of doxycycline and rifampin for patients with Alzheimer's disease. *J Am Geriatr Soc* 2004 March;52(3):381-7.
- (473) Molloy DW, Standish TI, Zhou Q, Guyatt G. A multicenter, blinded, randomized, factorial controlled trial of doxycycline and rifampin for treatment of Alzheimer's disease: the DARAD trial. *Int J Geriatr Psychiatry* 2012 June 21.
- (474) Lee CY, Landreth GE. The role of microglia in amyloid clearance from the AD brain. *J Neural Transm* 2010 August;117(8):949-60.
- (475) Terwel D, Steffensen KR, Verghese PB et al. Critical role of astroglial apolipoprotein E and liver X receptor-alpha expression for microglial Abeta phagocytosis. *J Neurosci* 2011 May 11;31(19):7049-59.
- (476) Lee CY, Tse W, Smith JD, Landreth GE. Apolipoprotein E promotes beta-amyloid trafficking and degradation by modulating microglial cholesterol levels. *J Biol Chem* 2012 January 13;287(3):2032-44.
- (477) Jiang Q, Lee CY, Mandrekar S et al. ApoE promotes the proteolytic degradation of Abeta. *Neuron* 2008 June 12;58(5):681-93.
- (478) Riddell DR, Zhou H, Comery TA et al. The LXR agonist TO901317 selectively lowers hippocampal Abeta42 and improves memory in the Tg2576 mouse model of Alzheimer's disease. *Mol Cell Neurosci* 2007 April;34(4):621-8.
- (479) Fitz NF, Cronican A, Pham T et al. Liver X receptor agonist treatment ameliorates amyloid pathology and memory deficits caused by high-fat diet in APP23 mice. *J Neurosci* 2010 May 19;30(20):6862-72.
- (480) Cui W, Sun Y, Wang Z, Xu C, Peng Y, Li R. Liver X receptor activation attenuates inflammatory response and protects cholinergic neurons in APP/PS1 transgenic mice. *Neuroscience* 2012 May 17;210:200-10.

- (481) Hoang MH, Jia Y, Jun HJ et al. Taurine is a liver X receptor-alpha ligand and activates transcription of key genes in the reverse cholesterol transport without inducing hepatic lipogenesis. *Mol Nutr Food Res* 2012 June;56(6):900-11.
- (482) Cramer PE, Cirrito JR, Wesson DW et al. ApoE-directed therapeutics rapidly clear beta-amyloid and reverse deficits in AD mouse models. *Science* 2012 March 23;335(6075):1503-6.
- (483) Mandrekar-Colucci S, Karlo JC, Landreth GE. Mechanisms Underlying the Rapid Peroxisome Proliferator-Activated Receptor-gamma-Mediated Amyloid Clearance and Reversal of Cognitive Deficits in a Murine Model of Alzheimer's Disease. *J Neurosci* 2012 July 25;32(30):10117-28.
- (484) Searcy JL, Phelps JT, Pancani T et al. Long-term pioglitazone treatment improves learning and attenuates pathological markers in a mouse model of Alzheimer's disease. *J Alzheimers Dis* 2012;30(4):943-61.
- (485) Jiang LY, Tang SS, Wang XY et al. PPARgamma Agonist Pioglitazone Reverses Memory Impairment and Biochemical Changes in a Mouse Model of Type 2 Diabetes Mellitus. *CNS Neurosci Ther* 2012 August;18(8):659-66.
- (486) Escribano L, Simon AM, Gimeno E et al. Rosiglitazone rescues memory impairment in Alzheimer's transgenic mice: mechanisms involving a reduced amyloid and tau pathology. *Neuropsychopharmacology* 2010 June;35(7):1593-604.
- (487) O'Reilly JA, Lynch M. Rosiglitazone improves spatial memory and decreases insoluble Abeta(1-42) in APP/PS1 mice. *J Neuroimmune Pharmacol* 2012 March;7(1):140-4.
- (488) Miller BW, Willett KC, Desilets AR. Rosiglitazone and pioglitazone for the treatment of Alzheimer's disease. *Ann Pharmacother* 2011 November;45(11):1416-24.
- (489) Nissen SE, Wolski K. Effect of rosiglitazone on the risk of myocardial infarction and death from cardiovascular causes. *N Engl J Med* 2007 June 14;356(24):2457-71.
- (490) Singh S, Loke YK, Furberg CD. Long-term risk of cardiovascular events with rosiglitazone: a meta-analysis. *JAMA* 2007 September 12;298(10):1189-95.
- (491) Gilman S, Koller M, Black RS et al. Clinical effects of Abeta immunization (AN1792) in patients with AD in an interrupted trial. *Neurology* 2005 May 10;64(9):1553-62.
- (492) Green RC, Schneider LS, Amato DA et al. Effect of tarenflurbil on cognitive decline and activities of daily living in patients with mild Alzheimer disease: a randomized controlled trial. *JAMA* 2009 December 16;302(23):2557-64.
- (493) Korczyn AD. Why have we failed to cure Alzheimer's disease? *J Alzheimers Dis* 2012;29(2):275-82.
- (494) Shi YQ, Huang TW, Chen LM et al. Ginsenoside Rg1 attenuates amyloid-beta content, regulates PKA/CREB activity, and improves cognitive performance in SAMP8 mice. *J Alzheimers Dis* 2010;19(3):977-89.
- (495) Janssen IM, Sturtz S, Skipka G, Zentner A, Velasco GM, Busse R. Ginkgo biloba in Alzheimer's disease: a systematic review. *Wien Med Wochenschr* 2010 December;160(21-22):539-46.

- (496) Jayaprakasam B, Padmanabhan K, Nair MG. Withanamides in *Withania somnifera* fruit protect PC-12 cells from beta-amyloid responsible for Alzheimer's disease. *Phytother Res* 2010 June;24(6):859-63.
- (497) Pase MP, Kean J, Sarris J, Neale C, Scholey AB, Stough C. The Cognitive-Enhancing Effects of *Bacopa monnieri*: A Systematic Review of Randomized, Controlled Human Clinical Trials. *J Altern Complement Med* 2012 July;18(7):647-52.